

Datasheet for ABIN3029328
anti-Tuberin antibody (pSer1798)



[Go to Product page](#)

3 Images

Overview

Quantity:	0.4 mL
Target:	Tuberin (TSC2)
Binding Specificity:	pSer1798
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tuberin antibody is un-conjugated
Application:	ELISA, Dot Blot (DB), Immunofluorescence (IF)

Product Details

Immunogen:	This phospho-TSC2 antibody was produced from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding pS1798 of human TSC2.
Isotype:	Ig Fraction
Purification:	Antigen affinity purified

Target Details

Target:	Tuberin (TSC2)
Alternative Name:	TSC2 (TSC2 Products)
Background:	Mutations in TSC2 lead to tuberous sclerosis complex. The protein is believed to be a tumor suppressor and is able to specifically stimulate the intrinsic GTPase activity of the Ras-related

Target Details

protein RAP1A and RAB5. The protein associates with hamartin in a cytosolic complex, possibly acting as a chaperone for hamartin. TSC2 may have a function in vesicular transport, but may also play a role in the regulation of cell growth arrest and in the regulation of transcription mediated by steroid receptors. Interaction between TSC1 and TSC2 may facilitate vesicular docking.


UniProt:	P49815
Pathways:	RTK Signaling , AMPK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Regulation of Cell Size , Tube Formation , Protein targeting to Nucleus

Application Details

Application Notes:	Titration of the phospho-TSC2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.\. Immunofluorescence: 1:10-1:50, Dot blot: 1:500
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	In 1X PBS, pH 7.4, with 0.09 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Aliquot the phospho-TSC2 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.




Immunofluorescence

Image 1. Confocal immunofluorescent analysis of phospho-TSC2 antibody with MCF-7 cells followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 Phalloidin (red). DAPI was used as a nuclear counterstain



Dot Blot

Image 2. Dot blot analysis of phospho-TSC2 antibody. 50ng of phos-peptide or nonphos-peptide per dot were spotted.



Immunofluorescence

Image 3. Confocal immunofluorescent analysis of phospho-TSC2 antibody with MCF-7 cells followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 Phalloidin (red). DAPI was used as a nuclear counterstain (blue).