

Datasheet for ABIN3029394
anti-p53 antibody (AA 33-62)



[Go to Product page](#)

5 Images

Overview

Quantity:	0.4 mL
Target:	p53 (TP53)
Binding Specificity:	AA 33-62
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This p53 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	A portion of amino acids 33-62 from the human protein was used as the immunogen for this TP53 antibody.
Isotype:	Ig Fraction
Purification:	Antigen affinity purified

Target Details

Target:	p53 (TP53)
Alternative Name:	TP53 (p53) (TP53 Products)
Background:	Tumor protein p53, a nuclear protein, plays an essential role in the regulation of cell cycle, specifically in the transition from G0 to G1. It is found in very low levels in normal cells, however, in a variety of transformed cell lines, it is expressed in high amounts, and believed to contribute

Target Details

to transformation and malignancy. p53 is a DNA-binding protein containing DNA-binding, oligomerization and transcription activation domains. It is postulated to bind as a tetramer to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site, and hence cause the loss of tumor suppressor activity. Alterations of the TP53 gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome.


UniProt:	P04637
Pathways:	p53 Signaling , MAPK Signaling , PI3K-Akt Signaling , Apoptosis , AMPK Signaling , Chromatin Binding , ER-Nucleus Signaling , Positive Regulation of Endopeptidase Activity , Hepatitis C , Protein targeting to Nucleus , Autophagy , Warburg Effect

Application Details

Application Notes:	Titration of the TP53 antibody may be required due to differences in protocols and secondary/substrate sensitivity.\. Western blot: 1:1000,Immunofluorescence: 1:10-1:50,IHC (Paraffin): 1:10-1:50
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	In 1X PBS pH 7.4 with 0.09 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Aliquot the TP53 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.




Immunofluorescence

Image 1. Confocal immunofluorescent analysis of TP53 antibody with A2058 cells followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 Phalloidin (red). DAPI was used as a nuclear counterstain (bl



Immunohistochemistry

Image 2. TP53 antibody immunohistochemistry analysis in formalin fixed and paraffin embedded human skin tissue.



Western Blotting

Image 3. Western blot analysis of TP53 antibody and 293 cell lysate (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the TP53 gene (2).

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN3029394.