

## Datasheet for ABIN3029442

# anti-UHRF1 antibody (C-Term)

2 Images



Go to Product page

$\sim$			
	ve	r\/	٨
$\cup$	V C	1 V I	٧V

Overview			
Quantity:	0.4 mL		
Target:	UHRF1		
Binding Specificity:	C-Term		
Reactivity:	Mouse		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This UHRF1 antibody is un-conjugated		
Application:	Western Blotting (WB), ELISA		
Product Details			
Immunogen:	This mouse Uhrf1 antibody was produced from a rabbit immunized with a KLH conjugated		
Immunogen:	This mouse Uhrf1 antibody was produced from a rabbit immunized with a KLH conjugated synthetic peptide between 596-628 amino acids from the C-terminal region of mouse Uhrf1.		
Immunogen: Isotype:			
	synthetic peptide between 596-628 amino acids from the C-terminal region of mouse Uhrf1.		
Isotype: Purification:	synthetic peptide between 596-628 amino acids from the C-terminal region of mouse Uhrf1.  Ig Fraction		
Isotype: Purification: Target Details	synthetic peptide between 596-628 amino acids from the C-terminal region of mouse Uhrf1.  Ig Fraction  Antigen affinity purified		
Isotype: Purification:	synthetic peptide between 596-628 amino acids from the C-terminal region of mouse Uhrf1.  Ig Fraction		
Isotype: Purification: Target Details	synthetic peptide between 596-628 amino acids from the C-terminal region of mouse Uhrf1.  Ig Fraction  Antigen affinity purified		
Isotype: Purification: Target Details Target:	synthetic peptide between 596-628 amino acids from the C-terminal region of mouse Uhrf1.  Ig Fraction  Antigen affinity purified  UHRF1		
Isotype: Purification: Target Details Target: Alternative Name:	synthetic peptide between 596-628 amino acids from the C-terminal region of mouse Uhrf1.  Ig Fraction  Antigen affinity purified  UHRF1  Uhrf1 (UHRF1 Products)		
Isotype: Purification: Target Details Target: Alternative Name:	synthetic peptide between 596-628 amino acids from the C-terminal region of mouse Uhrf1.  Ig Fraction  Antigen affinity purified  UHRF1  Uhrf1 (UHRF1 Products)  Multidomain protein that acts as a key epigenetic regulator by bridging DNA methylation and		

of the DNA methylation patterns through DNA replication. In addition to its role in maintenance of DNA methylation, also plays a key role in chromatin modification: through its tudor-like regions and PHD- type zinc fingers, specifically recognizes and binds histone H3 trimethylated at 'Lys-9' (H3K9me3) and unmethylated at 'Arg-2' (H3R2me0), respectively, and recruits chromatin proteins. Enriched in pericentric heterochromatin where it recruits different chromatin modifiers required for this chromatin replication. Also localizes to euchromatic regions where it negatively regulates transcription possibly by impacting DNA methylation and histone modifications. Has E3 ubiquitin-protein ligase activity by mediating the ubiquitination of target proteins such as histone H3 and PML. It is still unclear how E3 ubiquitin-protein ligase activity is related to its role in chromatin in vivo. May be involved in DNA repair.

Titration of the Uhrf1 antibody may be required due to differences in protocols and


UniProt:


Q8VDF2

### **Application Details**

Application Notes:

	secondary/substrate sensitivity.\. Western blot: 1:1000-2000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	In 1X PBS, pH 7.4, with 0.09 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Aliquot the Uhrf1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw
	cycles.





#### **Western Blotting**

**Image 1.** Western blot testing of Uhrf1 antibody at 1:1000 dilution + 1) mouse heart and 2) mouse thymus lysate. Predicted molecular weight ~ 88 kDa.

#### **Western Blotting**

**Image 2.** Western blot testing of Uhrf1 antibody at 1:1000 dilution + 1) mouse heart and 2) mouse thymus lysate. Predicted molecular weight ~ 88 kDa.