

Datasheet for ABIN3029749

**anti-PIK3R1 antibody (AA 447-461)****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	PIK3R1 (PI3K p85a)
Binding Specificity:	AA 447-461
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PIK3R1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	Amino acids 447-461 (KVYHQYQDKSREYD-beta specific) was used as the immunogen for this PI3K antibody (100% human, mouse and rat homology).
Isotype:	IgG
Purification:	Antigen affinity

## Target Details

Target:	PIK3R1 (PI3K p85a)
Alternative Name:	PI3K ( <a href="#">PI3K p85a Products</a> )
Background:	Phosphatidylinositol 3-kinase regulatory subunit 2, also called p85-Beta, PIK3R2 and PIK3, is an enzyme that in humans is encoded by the PIK3R2 gene. The NS1 protein from various influenza strains bind p85-beta, but not p85-alpha. Binding of p85-beta and activation of PI3K required

## Target Details

tyr89 of NS1, and mutant viruses expressing NS1 with a tyr89-to-phe substitution grew more slowly in cell culture than wildtype viruses. Using mouse embryonic fibroblasts, Park et al. showed that, in addition to regulating PI3K function, p85-alpha/beta regulated the function of Xbp1s, a transcription factor that orchestrates the unfolded protein response (UPR) following endoplasmic reticulum (ER) stress. Overexpression of the protein in livers of ob/ob mice increased glucose tolerance and reduced blood glucose concentrations.

UniProt:	<a href="#">O00459</a>
Pathways:	<a href="#">TCR Signaling</a> , <a href="#">Response to Growth Hormone Stimulus</a> , <a href="#">Regulation of Muscle Cell Differentiation</a> , <a href="#">Skeletal Muscle Fiber Development</a> , <a href="#">Hepatitis C</a> , <a href="#">Protein targeting to Nucleus</a> , <a href="#">VEGF Signaling</a> , <a href="#">BCR Signaling</a> , <a href="#">Warburg Effect</a>


## Application Details

Application Notes:	The stated application concentrations are suggested starting points. Titration of the PI3K antibody may be required due to differences in protocols and secondary/substrate sensitivity.\. Western blot: 0.5-1 µg/mL,IHC (Paraffin): 0.5-1 µg/mL
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------


## Handling

Buffer:	0.5 mg/mL if reconstituted with 0.2 mL sterile DI water
Storage:	-20 °C
Storage Comment:	After reconstitution, the PI3K antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.




Immunohistochemistry

**Image 1.** IHC-P: PI3K antibody testing of human breast cancer tissue



Western Blotting

**Image 2.** Western blot testing of PI3K antibody and Lane 1: rat testis; 2: 293T; 3: HeLa cell lysate



Western Blotting

**Image 3.** Western blot testing of PI3K antibody and Lane 1: rat testis