# ANTIBODIES ONLINE

## Datasheet for ABIN3029987 anti-ATG7 antibody (AA 284-313)

5 Images



Overview


Quantity:	0.4 mL
Target:	ATG7
Binding Specificity:	AA 284-313
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATG7 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA
Product Details	
Immunogen:	A portion of amino acids 284-313 from the human protein was used as the immunogen for this
	APG7 antibody.
lsotype:	Ig Fraction
Cross-Reactivity (Details):	Expected species reactivity: Rat, Chicken
Purification:	Purified


### Target Details

Target:	ATG7
Alternative Name:	ATG7 (APG7) (ATG7 Products)
Background:	Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN3029987 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

	constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic
	enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of
	double-membrane bound autophagosomes which enclose the cytoplasmic constituent
	targeted for degradation in a membrane bound structure, which then fuse with the lysosome
	(or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded
	within the lysosome (or vacuole). APG7 functions as an E1 enzyme essential for
	multisubstrates such as GABARAPL1 and ATG12. APG3L is an E2-like conjugating enzyme
	facilitating covalent binding of APG8 (MAP1LC3) to phosphatidylethanolamine (PE). APG7 (an
	E1-like enzyme) facilitates this reaction by forming an E1-E2 complex with APG3. Formation of
	the PE conjugate is essential for autophagy.
UniProt:	095352
Pathways:	Response to Water Deprivation, Autophagy
Application Details	
	Titration of the APG7 antibody may be required due to differences in protocols and
Application Notes:	secondary/substrate sensitivity.\. Western blot: 1:1000,IHC (Paraffin): 1:50-1:100
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	In 1X PBS, pH 7.4, with 0.09 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Aliquot the APG7 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw
	cycles.





#### Immunohistochemistry

**Image 1.** IHC analysis of FFPE human breast carcinoma tissue stained with the APG7 antibody

#### Western Blotting

**Image 2.** Western blot analysis of APG7 antibody and 293 lysate transiently transfected with the ATG7 gene.

#### Western Blotting

**Image 3.** Western blot testing of APG7 antibody and mouse liver tissue lysate. Predicted molecular weight: 70-80 kDa.

Please check the product details page for more images. Overall 5 images are available for ABIN3029987.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN3029987 | 07/26/2024 | Copyright antibodies-online. All rights reserved.