

Datasheet for ABIN3030205
anti-BRD4 antibody (AA 1313-1342)



[Go to Product page](#)

3 Images

Overview

Quantity:	0.4 mL
Target:	BRD4
Binding Specificity:	AA 1313-1342
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BRD4 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	A portion of amino acids 1313-1342 from the human protein was used as the immunogen for this BRD4 antibody.
Isotype:	Ig Fraction
Cross-Reactivity (Details):	Expected species reactivity: Mouse
Purification:	Purified

Target Details

Target:	BRD4
Alternative Name:	BRD4 (BRD4 Products)
Background:	BRD4 is homologous to the murine protein MCAP, which associates with chromosomes during

Target Details

mitosis, and to the human RING3 protein, a serine/threonine kinase. Each of these proteins contains two bromodomains, a conserved sequence motif which may be involved in chromatin targeting. The gene has been implicated as the chromosome 19 target of translocation t(15,19)(q13,p13.1), which defines an upper respiratory tract carcinoma in young people.


UniProt:	O60885
Pathways:	Chromatin Binding , SARS-CoV-2 Protein Interactome

Application Details

Application Notes:	Titration of the BRD4 antibody may be required due to differences in protocols and secondary/substrate sensitivity.\. IHC (Paraffin): 1:50-1:100
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	In 1X PBS, pH 7.4, with 0.09 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Aliquot the BRD4 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.




Immunofluorescence

Image 1. Subnuclear distribution of cellular proteins. CHO Bgl40 cells grown on coverslips were either directly or after treatment with 0.5% Triton X-100, incubated with BRD4 antibody (left, center). Propidium iodide staining of cellular DNA (right).



Immunohistochemistry

Image 2. IHC analysis of FFPE human breast carcinoma tissue stained with the BRD4 antibody



Immunohistochemistry

Image 3. IHC analysis of FFPE human breast carcinoma tissue stained with the BRD4 antibody