antibodies -online.com





anti-CD163L1 antibody (AA 1401-1430)





Go to Product page

\sim			
	IV/E	۱/۱۲	$I \cap V$

Overview		
Quantity:	0.4 mL	
Target:	CD163L1	
Binding Specificity:	AA 1401-1430	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CD163L1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA	
Product Details		
Immunogen:	A portion of amino acids 1401-1430 from the human protein was used as the immunogen for	
	this CD163L1 antibody.	
Isotype:	lg Fraction	
Purification:	Antigen affinity purified	
Target Details		
Target:	CD163L1	
Alternative Name:	CD163L1 (CD163L1 Products)	
Background:	This gene encodes a member of the scavenger receptor cysteine-rich (SRCR) superfamily.	
	Members of this family are secreted or membrane-anchored proteins mainly found in cells	
	associated with the immune system. The SRCR family is defined by a 100-110 amino acid	

Target Details

SRCR domain, which may mediate protein-protein interaction and ligand binding. The encoded protein contains twelve SRCR domains, a transmembrane region and a cytoplasmic domain. Alternatively spliced transcript variants encoding different isoforms have been described but their full-length nature has not been determined.

UniProt:

Q9NR16


Application Details

Application Notes:	Titration of the CD163L1 antibody may be required due to differences in protocols and	
	secondary/substrate sensitivity.\. Western blot: 1:1000	
Restrictions:	For Research Use only	

Handling


Format:	Liquid	
Buffer:	In 1X PBS pH 7.4 with 0.09 % sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Aliquot the CD163L1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.	

Images



Western Blotting

Image 1. CD163L1 antibody western blot analysis in NCI-H292 lysate



Western Blotting

Image 2. CD163L1 antibody western blot analysis in NCI-H292 lysate