

Datasheet for ABIN3030641 anti-CDKN2A antibody (pSer8)

2 Images



Go to Product page

| _ | | | | |
|-----|-----|-----|-----|---|
| () | ve. | rv/ | 101 | Λ |

| Quantity: | 0.4 mL | |
|----------------------|--|--|
| Target: | CDKN2A | |
| Binding Specificity: | pSer8 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This CDKN2A antibody is un-conjugated | |
| Application: | ELISA, Dot Blot (DB) | |
| Product Details | | |
| Immunogen: | This phospho-CDKN2A antibody was produced from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding pS8 of human CDKN2A. | |
| Isotype: | Ig Fraction | |
| Purification: | Antigen affinity purified | |
| Target Details | | |
| Target: | CDKN2A | |
| Alternative Name: | CDKN2A (CDKN2A Products) | |
| Background: | This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which | |
| | | |

encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene, this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, MDM1, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene.

UniProt:

P42771

Pathways:

Mitotic G1-G1/S Phases, Stem Cell Maintenance, Positive Regulation of Endopeptidase Activity, Autophagy, Positive Regulation of Response to DNA Damage Stimulus

Application Details

Application Notes:


Titration of the phospho-CDKN2A antibody may be required due to differences in protocols and secondary/substrate sensitivity.\. Dot blot: 1:500

Restrictions:

For Research Use only


Handling

| Format: | Liquid | |
|--------------------|--|--|
| Buffer: | In 1X PBS, pH 7.4, with 0.09 % sodium azide | |
| Preservative: | Sodium azide | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Aliquot the phospho-CDKN2A antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles. | |



Dot Blot

Image 1. Dot blot analysis of phospho-CDKN2A antibody. 50ng of phos-peptide or nonphos-peptide per dot were spotted.



Dot Blot

Image 2. Dot blot analysis of phospho-CDKN2A antibody. 50ng of phos-peptide or nonphos-peptide per dot were spotted.