antibodies - online.com







anti-Desmoglein 3 antibody (AA 981-999)

Images



Go to	Droc	LIOT	nagg
GU LU	FIUU	IUCL	paut

()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	1 V I	ew

Overview	
Quantity:	100 μg
Target:	Desmoglein 3 (DSG3)
Binding Specificity:	AA 981-999
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Desmoglein 3 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	Amino acids 981-999 (QLRGSHTMLCTEDPCSRLI-human) were used as the immunogen for
	this Desmoglein 3 antibody.
Isotype:	IgG
Purification:	Antigen affinity
Target Details	
Target:	Desmoglein 3 (DSG3)
Alternative Name:	Desmoglein 3 (DSG3 Products)
Background:	Desmoglein 3 is also known as PVA (Pemphigus Vulgaris Antigen). Pemphigus vulgaris (PV)
	and pemphigus foliaceus (PF) are autoimmune diseases of the skin which have as target
	cutions a true different respective of the decreased in subfamily of the decrease and called in a
	antigens two different members of the desmoglein subfamily of the desmosomal cadherins:

Target Details

pemphigus vulgaris antigen (PVA, or Desmoglein 3) in the case of PV and Desmoglein 1 in the case of PF. The DSG3 gene, like the genes for DSG1 and DSG2, is located on chromosome 18. The human DSG3 gene consists of 15 exons and spans more than 23 kb. DSG3 and DSG1 are the autoantigens of pemphigus vulgaris and pemphigus foliaceus, respectively. DSG3 and DSG1 are the target antigens in PNP, and IgG autoantibodies against DSG3 in PNP sera play a pathogenic role in the loss of keratinocyte adhesion and blistering.

UniProt:

P32926

Application Details

Application Notes:	The stated application concentrations are suggested starting amounts. Titration of the
	Desmoglein 3 antibody may be required due to differences in protocols and
	secondary/substrate sensitivity.\. Western blot: 0.5-1 µg/mL

Restrictions:

For Research Use only

Handling

Buffer:	0.5 mg/mL if reconstituted with 0.2 mL sterile DI water
Storage:	-20 °C
Storage Comment: After reconstitution, the Desmoglein 3 antibody can be stored for up to one month at 4° long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing	

Images



Western Blotting

Image 1. Western blot testing of Desmoglein 3 antibody and Lane 1: A431



Western Blotting

Image 2. Western blot testing of Desmoglein 3 antibody and

Lane 1: A431; 2: CEM