

# Datasheet for ABIN3030827

# anti-EPHA1 antibody (N-Term)

# **Images**



_			
	Ve.	rv	iew

Quantity:	0.4 mL	
Target:	EPHA1	
Binding Specificity:	AA 70-105, N-Term	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This EPHA1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA	
Product Details		
Immunogen:	This mouse Epha1 antibody was produced from a rabbit immunized with a KLH conjugated	
Immunogen:	This mouse Epha1 antibody was produced from a rabbit immunized with a KLH conjugated synthetic peptide between 70-105 amino acids from the N-terminal region of mouse Epha1.	
Immunogen: Isotype:		
	synthetic peptide between 70-105 amino acids from the N-terminal region of mouse Epha1.	
Isotype: Purification:	synthetic peptide between 70-105 amino acids from the N-terminal region of mouse Epha1.  Ig Fraction	
Isotype:	synthetic peptide between 70-105 amino acids from the N-terminal region of mouse Epha1.  Ig Fraction	
Isotype: Purification:	synthetic peptide between 70-105 amino acids from the N-terminal region of mouse Epha1.  Ig Fraction	
Isotype: Purification: Target Details	synthetic peptide between 70-105 amino acids from the N-terminal region of mouse Epha1.  Ig Fraction  Antigen affinity purified	
Isotype: Purification: Target Details Target:	synthetic peptide between 70-105 amino acids from the N-terminal region of mouse Epha1.  Ig Fraction  Antigen affinity purified  EPHA1	
Isotype: Purification: Target Details Target: Alternative Name:	synthetic peptide between 70-105 amino acids from the N-terminal region of mouse Epha1.  Ig Fraction  Antigen affinity purified  EPHA1  Epha1 (EPHA1 Products)	

### **Target Details**

while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Binds with a low affinity EFNA3 and EFNA4 and with a high affinity to EFNA1 which most probably constitutes its cognate/functional ligand. Upon activation by EFNA1 induces cell attachment to the extracellular matrix inhibiting cell spreading and motility through regulation of ILK and downstream RHOA and RAC. Plays also a role in angiogenesis and regulates cell proliferation. May play a role in apoptosis.

UniProt:


Q60750

## Application Details

Application Notes:	Titration of the Epha1 antibody may be required due to differences in protocols and		
	secondary/substrate sensitivity.\. Western blot: 1:1000		
Restrictions:	For Research Use only		
Handling			

#### Handling

Format:	Liquid	
Buffer:	In 1X PBS, pH 7.4, with 0.09 % sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Aliquot the Epha1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.	



### **Western Blotting**

**Image 1.** Western blot analysis of lysate from mouse 1) kidney, 2) lung and 3) skin tissue using Epha1 antibody at 1:1000.

### **Western Blotting**

**Image 2.** Western blot analysis of lysate from mouse 1) kidney, 2) lung and 3) skin tissue using Epha1 antibody at 1:1000.