

Datasheet for ABIN3031839
anti-MEK1 antibody (C-Term)



[Go to Product page](#)

6 Images

Overview

Quantity:	100 µg
Target:	MEK1 (MAP2K1)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MEK1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

Product Details

Immunogen:	An amino acid sequence from the C-terminus of human MEK1 (KQLMVHAFIKRSDAE) was used as the immunogen for this MEK1 antibody (100% homologous in human, mouse and rat).
Isotype:	IgG
Purification:	Antigen affinity

Target Details

Target:	MEK1 (MAP2K1)
Alternative Name:	MEK1 (MAP2K1 Products)
Background:	Dual specificity mitogen-activated protein kinase kinase 1 is an enzyme that in humans is encoded by the MAP2K1 gene. The protein encoded by this gene is a member of the dual

Target Details

specificity protein kinase family, which acts as a mitogen-activated protein(MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases(ERKs), act as an integration point for multiple biochemical signals. MEK1 lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon activation by a wide variety of extra- and intracellular signals. As an essential component of the MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development. Rampoldi et al.(1997) localized the MAP2K1 gene to 15q22.1-q22.33.


UniProt:	Q02750
Pathways:	MAPK Signaling , RTK Signaling , Interferon-gamma Pathway , Fc-epsilon Receptor Signaling Pathway , Neurotrophin Signaling Pathway , Activation of Innate immune Response , Toll-Like Receptors Cascades , Autophagy , Signaling of Hepatocyte Growth Factor Receptor , BCR Signaling

Application Details

Application Notes:	The stated application concentrations are suggested starting amounts. Titration of the MEK1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.\. Western blot: 0.5-1 µg/mL,IHC (Paraffin): 0.5-1 µg/mL,Immunocytochemistry: 0.5-1 µg/mL
Restrictions:	For Research Use only


Handling

Buffer:	0.5 mg/mL if reconstituted with 0.2 mL sterile DI water
Storage:	-20 °C
Storage Comment:	After reconstitution, the MEK1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.




Western Blotting

Image 1. Western blot testing of MEK1 antibody and Lane 1: rat skeletal muscle



Immunohistochemistry

Image 2. IHC-P: MEK1 antibody testing of human breast cancer tissue



Immunohistochemistry

Image 3. IHC-P: MEK1 antibody testing of human lung cancer tissue

Please check the [product details page](#) for more images. Overall 6 images are available for ABIN3031839.