antibodies -online.com







anti-MUC4 antibody (AA 1249-1444)

Images



| \sim | |
|--------|----------|
| ()\/\ | rview |
| \cup | 1 410 44 |

Background:

| Quantity: | 100 μg |
|----------------------|---|
| Target: | MUC4 |
| Binding Specificity: | AA 1249-1444 |
| Reactivity: | Human, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MUC4 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | Human partial recombinant protein (1249-1444) was used as the immunogen for this MUC4 |
| | antibody. |
| Isotype: | lg Fraction |
| Purification: | Antigen affinity |
| Target Details | |
| Target: | NUO 4 |
| rarget. | MUC4 |

Mucin 4, also known as MUC4, is a mucin protein that in humans is encoded by the MUC4

In normal functioning, Mucin 4 is known to play anti-adhesive roles in the body, such as in

gene. Like other mucins, Mucin 4 is a high-molecular weight glycoprotein. It is mapped to 3q29.

Target Details

lubricating the reproductive lining. It is found in the respiratory tract, particularly in the trachea and lung and the digestive tract, in the esophagus and colon, as well as in the visual and auditory systems. Mucin 4 serves to protect and lubricate the epithelium, which facilitates transport and traps foreign particles. It is found to be overexpressed during the luteal phase of menstruation. Mucin 4 may also have a role in fetal morphogenic development.

Gene ID:

4585

Pathways:

Regulation of Leukocyte Mediated Immunity

Application Details

Application Notes:

The stated application concentrations are suggested starting points. Titration of the MUC4 antibody may be required due to differences in protocols and secondary/substrate sensitivity.\. Western blot: $0.5-1~\mu g/mL$

Restrictions:

For Research Use only

Handling

Buffer:

0.5 mg/mL if reconstituted with 0.2 mL sterile DI water

Storage:

-20 °C

Storage Comment:

After reconstitution, the MUC4 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

Images



6-

Western Blotting

Image 1. Western blot testing of MUC4 antibody and PC12 lysate

kDa 200 - 116 - 97 - 66 - 44 - 31 - 22 - 14 - 6 -

Western Blotting

Image 2. Western blot testing of MUC4 antibody and PC12 lysate