# antibodies -online.com





### anti-AQP6 antibody (C-Term)





#### Overview

- Overview	
Quantity:	100 μg
Target:	AQP6
Binding Specificity:	AA 264-282, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AQP6 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Purpose:	Rabbit IgG polyclonal antibody for Aquaporin-6(AQP6) detection. Tested with WB, IHC-P in Human.
Immunogen:	A synthetic peptide corresponding to a sequence at the C-terminus of human Aquaporin 6(264-282aa EPLKKESQPGSGAVEMESV).
Sequence:	EPLKKESQPG SGAVEMESV
Isotype:	IgG
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Characteristics:	Rabbit IgG polyclonal antibody for Aquaporin-6(AQP6) detection. Tested with WB, IHC-P in Human.  Gene Name: aquaporin 6, kidney specific  Protein Name: Aquaporin-6

## **Product Details** Purification: Immunogen affinity purified. **Target Details** Target: AQP6 AQP6 (AQP6 Products) Alternative Name Background: Aquaporin 6, kidney specific is a protein in humans that is encoded by the AQP6 gene. The aquaporins(AQPs) are a family of water-transporting proteins that facilitate osmotically driven water movement across cell plasma membranes. Among the seven human aquaporins cloned to date(AQPs 0-6), genes encoding the four most closely related aquaporins(AQP0, AQP2, AQP5, and AQP6) have been mapped to chromosome band 12q13, suggesting an aquaporin family gene cluster at this locus. Unlike other aquaporins, AQP6 functions not as a water channel but as an anion-selective channel. Single-channel analyses have shown AQP6 to flicker rapidly between closed and open status. Synonyms: AQP-6 antibody|AQP2L antibody|Aqp6 antibody|AQP6\_HUMAN antibody|Aquaporin 2 like antibody|aquaporin 2 like kidney specific antibody|aquaporin 2-like kidney specific antibody|Aquaporin 6 antibody|Aquaporin 6 kidney specific antibody|Aquaporin-2-like antibody|Aquaporin-6 antibody|Aquaporin-6 kidney specific antibody|hKID antibody|Human kidney water channel(hKID) mRNA, complete cds antibody|KID antibody|Kidney-specific aquaporin antibody UniProt: Q13520 **Application Details**

Application Notes:	WB: Concentration: 0.1-0.5 μg/mL, Tested Species: Human
	IHC-P: Concentration: 0.5-1 μg/mL, Tested Species: Human, Epitope Retrieval by Heat: Boiling
	the paraffin sections in 10 mM citrate buffer, pH 6.0, for 20 mins is required for the staining of
	formalin/paraffin sections.
	Notes: Tested Species: Species with positive results. Predicted Species: Species predicted to be
	fit for the product based on sequence similarities. Other applications have not been tested.
	Optimal dilutions should be determined by end users.

Comment:

Antibody can be supported by chemiluminescence kit ABIN921124 in WB, supported by ABIN921231 in IHC(P).


Restrictions:

For Research Use only

#### Handling


Format:	Lyophilized
Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 μg/mL
Buffer:	Each vial contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na2HPO4, 0.05 mg Thimerosal, 0.05 mg Sodium azide.
Preservative:	Thimerosal (Merthiolate), Sodium azide
Precaution of Use:	This product contains Sodium azide and Thimerosal (Merthiolate): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	At -20°C for one year. After reconstitution, at 4°C for one month.  It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Expiry Date:	12 months

#### **Images**




#### Immunohistochemistry

**Image 1.** Anti-Aquaporin 6 antibody, IHC(P) IHC(P): Human Placenta Tissue



#### Immunohistochemistry

Image 2. Anti-Aquaporin 6 antibody, IHC(P) IHC(P): Human Lung Cancer Tissue



#### **Western Blotting**

Image 3. Anti-Aquaporin 6 antibody, Western blotting Lane1: U87 Cell Lysate Lane 2: COLO320 Cell Lysate Lane 3: MCF-7 Cell Lysate