antibodies.com

# Datasheet for ABIN3043427 anti-EIF6 antibody (AA 66-210)

6 Images



#### Overview

Quantity:	100 µg					
Target:	EIF6					
Binding Specificity:	AA 66-210					
Reactivity:	Human, Mouse, Rat					
Host:	Rabbit					
Clonality:	Polyclonal					
Conjugate:	This EIF6 antibody is un-conjugated					
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))					
Product Details						
Purpose:	Rabbit IgG polyclonal antibody for Eukaryotic translation initiation factor 6(EIF6) detection.					
	Tested with WB, IHC-P in Human,Mouse, Rat.					
Immunogen:	E.coli-derived human EIF6 recombinant protein (Position: N66-T210). Human EIF6 shares					
	99.3% amino acid (aa) sequence identity with both mouse and rat EIF6.					
lsotype:	lgG					
Cross-Reactivity (Details):	No cross reactivity with other proteins.					
Characteristics:	Rabbit IgG polyclonal antibody for Eukaryotic translation initiation factor 6(EIF6) detection.					
	Tested with WB, IHC-P in Human,Mouse, Rat.					
	Gene Name: eukaryotic translation initiation factor 6					
	Protein Name: Eukaryotic translation initiation factor 6					
Purification:	Immunogen affinity purified.					

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN3043427 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

## Target Details

Target:	EIF6				
Alternative Name:	EIF6 (EIF6 Products)				
Background:	EIF6 (Eukaryotic Translation Initiation Factor 6), also called EIF3A or ITGB4BP, is a human gene				
	By fluorescence in situ hybridization, Sanvito et al. (1998) mapped the ITGB4BP gene to				
	20q11.2. Ceci et al. (2003) demonstrated that the ribosomal 60S subunit is activated by release				
	of EIF6. In the cytoplasm, EIF6 is bound to free 60S but not to 80S subunits. Furthermore, EIF6				
	interacts in the cytoplasm with RACK1, a receptor for activated protein kinase C. Gandin et al.				
	(2008) demonstrated that mammalian eIF6 is required for efficient initiation of translation in				
	vivo. Eif6-null mouse embryos were lethal at preimplantation. Heterozygous mice had 50 %				
	reduction of eIF6 levels in all tissues, and showed reduced mass of hepatic and adipose tissues				
	due to a lower number of cells and to impaired G1/S cell cycle progression.				
	Synonyms: b(2)gcn antibody B(2)GCN homolog antibody B4 integrin interactor antibody Binding				
	protein of beta-4 integrin antibody CAB antibody  eIF-6 antibody EIF3A antibody EIF6				
	antibody Eukaryotic translation initiation factor 3A antibody Eukaryotic translation initiation				
	factor 6 antibody IF6_HUMAN antibody ITGB4BP antibody OK/SW-cl.27 antibody p27 beta 4				
	integrin binding protein antibody p27(BBP) antibody p27BBP antibody RP4-61404.1 antibody				
Gene ID:	3692				
UniProt:	P56537				
Pathways:	Ribonucleoprotein Complex Subunit Organization, Ribosome Assembly				
Application Details					
Application Notes:	WB: Concentration: 0.1-0.5 µg/mL, Tested Species: Human, Mouse, Rat				
	IHC-P: Concentration: 0.5-1 $\mu$ g/mL, Tested Species: Human, Mouse, Rat, Epitope Retrieval by				
	Heat: Boiling the paraffin sections in 10 mM citrate buffer, pH 6.0, for 20 mins is required for the				
	staining of formalin/paraffin sections.				
	Notes: Tested Species: Species with positive results. Other applications have not been tested.				
	Optimal dilutions should be determined by end users.				
Comment:	Antibody can be supported by chemiluminescence kit ABIN921124 in WB, supported by				
	ABIN921231 in IHC(P).				
Restrictions:	For Research Use only				


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3043427 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

## Handling

Format:	Lyophilized					
Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 $\mu$ g/mL.					
Concentration:	500 μg/mL					
Buffer:	Each vial contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na2HPO4, 0.05 mg Sodium azide.					
Preservative:	Sodium azide					
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.					
Handling Advice:	Avoid repeated freezing and thawing.					
Storage:	4 °C/-20 °C					
Storage Comment:	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.					

### Images

130KD-1 2 3 4 5 6	Western Blotting
100KD-	Image 1.
70KD -	
55KD —	
35KD-	
25KD	
15KD -	



Immunohistochemistry									
Image	2.	Anti-	EIF6	Picoband	antibody,IHC(P)	IHC(P):			
Mouse Intestine Tissue									

#### Immunohistochemistry

**Image 3.** Anti- EIF6 Picoband antibody,IHC(P) IHC(P): Human Intestinal Cancer Tissue

Please check the product details page for more images. Overall 6 images are available for ABIN3043427.