

Datasheet for ABIN3043867

anti-KCNMA1 antibody (AA 124-467)



[Go to Product page](#)

4 Images

Overview

Quantity:	100 µg
Target:	KCNMA1
Binding Specificity:	AA 124-467
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCNMA1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Purpose:	Rabbit IgG polyclonal antibody for Calcium-activated potassium channel subunit alpha-1(KCNMA1) detection. Tested with WB, IHC-P in Human,Mouse,Rat.
Immunogen:	E.coli-derived human KCNMA1 recombinant protein (Position: K124-Q467). Human KCNMA1 shares 99% amino acid (aa) sequence identity with both mouse and rat KCNMA1.
Isotype:	IgG
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Characteristics:	<p>Rabbit IgG polyclonal antibody for Calcium-activated potassium channel subunit alpha-1(KCNMA1) detection. Tested with WB, IHC-P in Human,Mouse,Rat.</p> <p>Gene Name: potassium large conductance calcium-activated channel, subfamily M, alpha member 1</p> <p>Protein Name: Calcium-activated potassium channel subunit alpha-1</p>

Product Details

Purification: Immunogen affinity purified.

Target Details

Target: KCNMA1

Alternative Name: KCNMA1 ([KCNMA1 Products](#))

Background: Calcium-activated potassium channel subunit alpha-1 also known as KCa1.1, or BK channel for short, is a voltage gated potassium channel encoded by the KCNMA1 gene and characterized by their large conductance of potassium ions (K⁺) through cell membranes. This gene is located on 10q22.3. BK channels are activated (opened) by changes in membrane electrical potential and/or by increases in concentration of intracellular calcium ion (Ca²⁺). It is essential for the regulation of several key physiological processes including smooth muscle tone and neuronal excitability. BK channels also contribute to the behavioral effects of ethanol in the worm *C. elegans* under high concentrations (> 100 mM, or approximately 0.50 % BAC).It remains to be determined if BK channels contribute to intoxication in humans.

Synonyms: subfamily M subunit alpha-1 antibody|bA205K10.1 antibody|BK channel antibody|BKCA alpha antibody|BKCA alpha subunit antibody|BKTM antibody|Calcium activated potassium channel subfamily M subunit alpha 1 antibody|Calcium activated potassium channel subunit alpha 1 antibody|Calcium-activated potassium channel antibody|Calcium-activated potassium channel subunit alpha-1 antibody|DKFZp686K1437 antibody|Drosophila slowpoke like antibody|hSlo antibody|K(VCA)alpha antibody|KCa1.1 antibody|KCMA1_HUMAN antibody|KCNMA 1 antibody|KCNMA antibody|KCNMA1 antibody|Large conductance calcium activated potassium channel subfamily M alpha member 1 antibody|Maxi K antibody|Maxi K channel antibody|Maxi Potassium channel alpha antibody|MaxiK antibody|MGC71881 antibody|OTTHUMP00000060198 antibody|OTTHUMP00000064154 antibody|OTTHUMP00000064155 antibody|OTTHUMP00000064156 antibody|OTTHUMP00000064157 antibody|OTTHUMP00000064158 antibody|OTTHUMP00000064159 antibody|OTTHUMP00000064160 antibody|OTTHUMP00000064161 antibody|OTTHUMP00000064162 antibody|OTTHUMP00000064164 antibody|OTTHUMP00000064165 antibody|Potassium large conductance calcium activated channel subfamily M alpha member 1 antibody|SAKCA antibody|Slo 1 antibody|SLO alpha antibody|SLO antibody|Slo homolog antibody|Slo-alpha antibody|Slo1 antibody|Slowpoke homolog antibody|Stretch activated Kca channel antibody

Gene ID: 3778

Target Details

UniProt: [Q12791](#)

Pathways: [Regulation of Hormone Metabolic Process, Sensory Perception of Sound](#)

Application Details

Application Notes: WB: Concentration: 0.1-0.5 µg/mL, Tested Species: Mouse, Rat, Predicted Species: Human, The detection limit for KCNMA1 is approximately 0.25 ng/lane under reducing conditions.
IHC-P: Concentration: 0.5-1 µg/mL, Tested Species: Human, Mouse, Rat, Epitope Retrieval by Heat: Boiling the paraffin sections in 10 mM citrate buffer, pH 6.0, for 20 mins is required for the staining of formalin/paraffin sections.
Notes: Tested Species: Species with positive results. Predicted Species: Species predicted to be fit for the product based on sequence similarities. Other applications have not been tested.
Optimal dilutions should be determined by end users.

Comment: Antibody can be supported by chemiluminescence kit ABIN921124 in WB, supported by ABIN921231 in IHC(P).

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.

Concentration: 500 µg/mL

Buffer: Each vial contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na₂HPO₄, 0.05 mg Sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Avoid repeated freezing and thawing.


Storage: 4 °C/-20 °C

Storage Comment: At -20°C for one year. After reconstitution, at 4°C for one month.
It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.




Western Blotting

Image 1. Anti- KCNMA1antibody, Western blotting All lanes: Anti KCNMA1 at 0.5ug/ml Lane 1: Rat Brain Tissue Lysate at 50ug Lane 2: Rat Testis Tissue Lysate at 50ug Lane 3: Mouse Brain Tissue Lysate at 50ug Lane 4: Mouse Testis Tissue Lysate at 50ug Predicted bind size: 137KD Observed bind size: 70KD



Immunohistochemistry

Image 2. Anti- KCNMA1antibody,IHC(P) IHC(P): Mouse Intestine Tissue



Immunohistochemistry

Image 3. Anti- KCNMA1antibody,IHC(P) IHC(P): Human Lung Cancer Tissue

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN3043867.