

[Go to Product page](#)

Datasheet for ABIN3044580 **anti-TCR beta antibody (FITC)**

Overview

Quantity:	100 µg
Target:	TCR beta
Reactivity:	Mouse
Host:	Armenian Hamster
Clonality:	Monoclonal
Conjugate:	This TCR beta antibody is conjugated to FITC
Application:	Flow Cytometry (FACS), Immunofluorescence (IF)

Product Details

Immunogen:	Affinity purified TCR from mouse DO-11.10 cells Species
Clone:	H57-597
Isotype:	IgG

Target Details

Target:	TCR beta
Alternative Name:	TCR beta (TCR beta Products)
Background:	T cell receptor (TCR) is a heterodimer consisting of an alpha and a beta chain (TCR alpha/beta) or a gamma and a delta chain (TCR gamma/delta). TCR-beta is a member of the immunoglobulin superfamily and a component of the CD3/TCR complex (along with TCR-alpha). It is expressed on alpha/beta TCR-bearing T cells and thymocytes. The CD3/TCR complex plays a key role in antigen recognition, signal transduction, and T cell activation.

Application Details

Application Notes:	It is recommended for use in flow cytometry. This reagent is effective for direct immunofluorescence staining of mouse tissue for flow cytometric analysis using 1 µg/10 ⁶ cells
Comment:	Fluorescein Isothiocyanate, FITC (Abs/Em: 495/ 517 nm).
Reagent Preparation:	1. Transfer the sample to a 12 x 75 mm polystyrene test tube (10 ⁶ cells). 2. Add anti-mouse anti-mouse TCR beta FITC and mix gently with a vortex mixer. The optimal volume should be determined by the individual laboratory. 3. The recommended negative control is a non-reactive FITC-conjugated antibody of the same isotype. 4. Incubate in the dark at room temperature at 4°C for 30 minutes or at room temperature (20-25 °C) for 15 minutes. 5. Add 1,5 mL of Lysing Solution to each sample and mix gently with a vortex mixer. Incubate for 10 minutes at room temperature in the dark. 6. Centrifuge at 1000 x g for 5 minutes. Gently aspirate the supernatant and discard it leaving approximately 50 µl of fluid. 7. Add 2 mL 0.01 mol/L PBS (It better that it containing 2% bovine serum albumin) and resuspend the cells by using a vortex mixer. 8. Centrifuge at 1000 x g for 5 minutes. Gently aspirate the supernatant and discard it leaving approximately 50 µl of fluid. 9. Resuspend pellet in an appropriate fluid for flow cytometry, e.g. 0.3 mL PBS. The PBS should contain 1% paraformaldehyde (fixative) if samples are not analysed the same day. 10. Analyse on a flow cytometer or store at 2-8 °C in the dark until analysis. Samples can be run up to 24 hours after lysis.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The conjugate is provided in liquid form in buffer containing 1% bovine serum albumin (BSA) and 0,09% Sodium azide, pH 7.2.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C