

Datasheet for ABIN3045753

Goat anti-Human IgA (Chain alpha) Antibody (TRITC)



Overview

Quantity:	0.5 mg
Target:	IgA
Binding Specificity:	Chain alpha
Reactivity:	Human
Host:	Goat
Conjugate:	TRITC
Application:	Flow Cytometry (FACS), Immunomicroscopy (IM)

Product Details

Immunogen:	Purified Human IgA, (alpha chain)
Fragment:	F(ab')2 fragment
Specificity:	Based on IEP, this antibody reacts with: heavy (α) chains on human IgA
Cross-Reactivity (Details):	Based on IEP, no reactivity is observed to: non-immunoglobulin human serum immunoglobulins light chains on all human immunoglobulins
Characteristics:	Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.
Purification:	Affinity purified using solid phase Human IgA
Purity:	> 90 % based on SDS-PAGE Small amounts of intact IgG may be present.

Target Details

rarget betails	
Target:	IgA
Abstract:	IgA Products
Target Type:	Antibody
Application Details	
Application Notes:	This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator.
Restrictions:	Suggested starting dilution(s): • 1:20 - 1:2,000 for most applications For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Rehydrate with 0.55 mL of deionized water and let stand 30 minutes at room temperature to dissolve. Centrifuge to remove any particulates. Prepare fresh working dilution daily.
Concentration:	1.0 mg/mL
Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free 0.05 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	2-8 °C Shelf Life: 1 year from date of receipt. Prepare working dilution prior to use and then discard.
Expiry Date:	12 months