

Datasheet for ABIN3045847

Goat anti-Human IgG (Fc Region) Antibody (DyLight 488)[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	IgG
Binding Specificity:	Fc Region
Reactivity:	Human
Host:	Goat
Conjugate:	DyLight 488
Application:	Flow Cytometry (FACS), Immunomicroscopy (IM)

Product Details

Immunogen:	Purified Human IgG, Fc fragment
Specificity:	Based on IEP, this antibody reacts with: heavy (γ) chains on human IgG
Cross-Reactivity (Details):	Based on IEP, no reactivity is observed to: non-immunoglobulin human serum immunoglobulins light chains on all human immunoglobulins
Characteristics:	Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.
Purification:	Affinity purified using solid phase Human IgG
Purity:	> 95 % based on SDS-PAGE

Target Details

Target: IgG

Abstract: [IgG Products](#)

Target Type: Antibody

Application Details

Application Notes: This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator.

Suggested starting dilution(s): • 1:20 - 1:2,000 for most applications

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Rehydrate with 1.1 mL of deionized water and let stand 30 minutes at room temperature to dissolve. Centrifuge to remove any particulates. Prepare fresh working dilution daily.

Concentration: 1.0 mg/mL

Buffer: 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free
0.05 % (w/v) Sodium Azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Storage Comment: Store freeze-dried powder at 2-8 °C.
Shelf Life: Product is stable for up to 4 weeks at 2-8°C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20°C.