

## Datasheet for ABIN3045938 Rabbit anti-Human IgG (Whole Molecule) Antibody (TRITC)



Overview	
Quantity:	1 mg
Target:	lgG
Binding Specificity:	Whole Molecule
Reactivity:	Human
Host:	Rabbit
Conjugate:	TRITC
Application:	Flow Cytometry (FACS), Immunomicroscopy (IM)
Product Details	
Immunogen:	Purified Human IgG, whole molecule
Specificity:	Based on IEP, this antibody reacts with: heavy (γ) chains on human IgG light chains on all human immunoglobulins
Cross-Reactivity (Details):	Based on IEP, no reactivity is observed to: non-immunoglobulin human serum immunoglobulins
Characteristics:	Rabbit serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.
Purification:	Affinity purified using solid phase Human IgG
Purity:	> 95 % based on SDS-PAGE

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN3045938 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Target:	lgG
Abstract:	IgG Products
Target Type:	Antibody
Application Details	
Application Notes:	This conjugate is suitable for immunomicroscopy and flow cytometry. The optimal working dilution should be determined by the investigator. Suggested starting dilution(s): • 1:20 - 1:2,000 for most applications
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Rehydrate with 1.1 mL of deionized water and let stand 30 minutes at room temperature to dissolve. Centrifuge to remove any particulates. Prepare fresh working dilution daily.
Concentration:	1.0 mg/mL
Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free 0.05 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	2-8 °C Shelf Life: 1 year from date of receipt. Prepare working dilution prior to use and then discard.
Expiry Date:	12 months