antibodies -online.com





anti-Fc gamma RII (CD32) antibody (Biotin)





Overview

| Quantity: | 100 μg |
|--------------|-----------------------|
| Target: | Fc gamma RII (CD32) |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | Biotin |
| Application: | Flow Cytometry (FACS) |

Product Details

| Immunogen: | K-562 human erythromyeloblastoid leukemia cell line |
|------------------|--|
| Clone: | IV-3 |
| Isotype: | IgG2b kappa |
| Characteristics: | IV.3 reacts with a low affinity receptor for the Fc of IgG (FcγRII, CD32) that is expressed on human platelets, monocytes, granulocytes and B cells. IV.3 also cross-reacts with granulocytes and monocytes from peripheral blood lymphocytes from non-human primates. |
| Purification: | Purified |
| Purity: | >95 % |
| Grade: | GMP Grade |

Target Details

| Target: | Fc gamma RII (CD32) |
|-------------------|--|
| Alternative Name: | CD32 (CD32 Products) |
| Background: | IV.3 reacts with a low affinity receptor for the Fc of IgG (FcyRII, CD32) that is expressed on human platelets, monocytes, granulocytes and B cells. IV.3 also cross-reacts with granulocytes and monocytes from peripheral blood lymphocytes from non-human primates. |
| NCBI Accession: | NM_021642 |
| UniProt: | P12318 |


Application Details

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 0.5 mg/mL |
| Buffer: | PBS pH 7.2, 0.09 % sodium azide |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C |

Images



Flow Cytometry

Image 1.