

Datasheet for ABIN3073961

TICAM2 Protein (AA 2-235) (His tag)



[Go to Product page](#)

1 Image

Overview

Quantity:	1 mg
Target:	TICAM2
Protein Characteristics:	AA 2-235
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This TICAM2 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence:	<p>GIGKSKINSC PLSLSWGKRH SVDTSPPGYHE SDSKKSEDLS LCNVAEHSNT TEGPTGKQEG</p> <p>AQSVEEMFEE EAEVEVFLKF VILHAEDDTD EALRVQNLLQ DDFGIKPGII FAEMPCGRQH</p> <p>LQNLDDAVNG SAWTILLTE NFLRDTWCNF QFYTSLMNSV NRQHKYNSVI PMRPLNNPLP</p> <p>RERTPFALQT INALEEESRG FPTQVERIFQ ESVYKTQQT WKETRNMVQR QFIA</p> <p>Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.</p>
-----------	---

Characteristics:	<ul style="list-style-type: none"> Made in Germany - from design to production - by highly experienced protein experts. Human TICAM2 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis).
------------------	--

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

Product Details

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the ExPASy's protparam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in bacterial culture: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Endotoxin has not been removed. Please contact us if you require endotoxin removal.
Grade:	Crystallography grade

Target Details

Target:	TICAM2
Alternative Name:	TICAM2 (TICAM2 Products)
Background:	Functions as sorting adapter in LPS-TLR4 signaling to regulate the MYD88-independent pathway during the innate immune response to LPS. Physically bridges TLR4 and TICAM1 and functionally transmits LPS-TLR4 signal to TICAM1, signaling is proposed to occur in early

Target Details

endosomes after endocytosis of TLR4. May also be involved in IL1-triggered NF-kappa-B activation, functioning upstream of IRAK1, IRAK2, TRAF6, and IKBKB, however, reports are controversial. Involved in IL-18 signaling and is proposed to function as a sorting adaptor for MYD88 in IL-18 signaling during adaptive immune response., Isoform 2: Proposed to inhibit LPS-TLR4 signaling at the late endosome by interaction with isoform 1 thereby disrupting the association of isoform 1 with TICAM1. May be involved in TLR4 degradation in late endosomes.

Molecular Weight: 27.7 kDa Including tag.

UniProt: [Q86XR7](#)

Pathways: [TLR Signaling](#), [Activation of Innate immune Response](#), [Cellular Response to Molecule of Bacterial Origin](#), [Toll-Like Receptors Cascades](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 100 mM NaCl, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: Unlimited (if stored properly)



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process