antibodies

# Datasheet for ABIN3074159 TGFB2 Protein (AA 303-414) (His tag)





#### Overview

Quantity:	1 mg
Target:	TGFB2
Protein Characteristics:	AA 303-414
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TGFB2 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)
Product Details	
Sequence:	ALDAAYCFRN VQDNCCLRPL YIDFKRDLGW KWIHEPKGYN ANFCAGACPY LWSSDTQHSR
	VLSLYNTINP EASASPCCVS QDLEPLTILY YIGKTPKIEQ LSNMIVKSCK CS
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a
	special request, please contact us.
Characteristics:	<ul> <li>Made in Germany - from design to production - by highly experienced protein experts.</li> <li>Human TGFB2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.</li> </ul>
	<ul> <li>State-of-the-art algorithm used for plasmid design (Gene synthesis).</li> </ul>
	This protein is a made to order protein and will be made for the first time for your order. Our
	experts in the lab will ensure that you receive a correctly folded protein.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN3074159 | 02/07/2024 | Copyright antibodies-online. All rights reserved.

(other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).         When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.         The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.         Purification:       Two step purification of proteins expressed in baculovirus infected SF9 insect cells:         1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.	Product Details	
(other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).         When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.         Purification:       Two step purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.         Purification:       2. Protein containing fractions of the best purification are subjected to second purification strony bize exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.         Purity:       >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.         Sterility:       0.22 µm filtered         Endotoxin Level:       Protein is endotoxin free.         Grade:       Crystallography grade		cannot be expressed or purified.
custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification si through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography grade		In the unlikely event that the protein cannot be expressed or purified we do not charge anything
experiments or purification optimization).         When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.         Purification:       Two step purification of proteins expressed in baculovirus infected SF9 insect cells: <ol> <li>In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li> <li>Protein containing fractions of the best purification are subjected to second purification step using size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE.</li> </ol> Purity:       >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.         Sterility:       0.22 µm filtered         Endotoxin Level:       Protein is endotoxin free.         Grade:       Crystallography grade		(other companies might charge you for any performed steps in the expression process for
When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm? The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.         Purification:       Two step purification of proteins expressed in baculovirus infected SF9 insect cells: <ul> <li>1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li> <li>2. Protein containing fractions of the best purification are subjected to second purification step, the protein is exclusion Chromatography and Western Blot.</li> </ul> Purity:       >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.         Sterility:       0.22 µm filtered         Endotoxin Level:       Protein is endotoxin free.         Grade:       Crystallography grade		custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression
Folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.         Purification:       Two step purification of proteins expressed in baculovirus infected SF9 insect cells: <ol> <li>In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li> <li>Protein containing fractions of the best purification are subjected to second purification step user blot.</li> </ol> <li>Purity: &gt;95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.</li> <li>Sterility: 0.22 µm filtered</li> <li>Endotoxin Level: Protein is endotoxin free.</li> <li>Grade: Crystallography grade</li> <li>Target Details</li>		experiments or purification optimization).
protein experts will do everything to make sure that you receive the protein you ordered.         The concentration of our recombinant proteins is measured using the absorbance at 280nm         The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.         The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.         Purification:       Two step purification of proteins expressed in baculovirus infected SF9 insect cells:         1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.         2. Protein containing fractions of the best purification are subjected to second purification step by SDS-PAGE.         2. Protein containing fractions of the best purification are analyzed by SDS-PAGE and Western Blot.         Purity:       >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.         Sterility:       0.22 µm filtered         Endotoxin Level:       Protein is endotoxin free.         Grade:       Crystallography grade         Target Details		When you order this made-to-order protein you will only pay upon receival of the correctly
The concentration of our recombinant proteins is measured using the absorbance at 280nm The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification st through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE 3. Purity:Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget Details		folded protein. With no financial risk on your end you can rest assured that our experienced
The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification si through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Endotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget Details		protein experts will do everything to make sure that you receive the protein you ordered.
specific reference buffer. The concentration of the protein is calculated using its specific absorption coefficient. We us the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification si through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget Details		The concentration of our recombinant proteins is measured using the absorbance at 280nm.
The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification st through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGEPurity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget Details		The protein's absorbance will be measured in several dilutions and is measured against its
the Expasy's protparam tool to determine the absorption coefficient of each protein.Purification:Two step purification of proteins expressed in baculovirus infected SF9 insect cells: 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification si through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget Details		specific reference buffer.
Purification:       Two step purification of proteins expressed in baculovirus infected SF9 insect cells:         1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.         2. Protein containing fractions of the best purification are subjected to second purification st through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.         Purity:       >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.         Sterility:       0.22 µm filtered         Endotoxin Level:       Protein is endotoxin free.         Grade:       Crystallography grade         Target Details		The concentration of the protein is calculated using its specific absorption coefficient. We use
<ul> <li>In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li> <li>Protein containing fractions of the best purification are subjected to second purification sit through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li> <li>Purity: &gt;95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.</li> <li>Sterility: 0.22 µm filtered</li> <li>Endotoxin Level: Protein is endotoxin free.</li> <li>Grade: Crystallography grade</li> <li>Target Details</li> </ul>		the Expasy's protparam tool to determine the absorption coefficient of each protein.
<ul> <li>different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li> <li>2. Protein containing fractions of the best purification are subjected to second purification sr through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li> <li>Purity: &gt;95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.</li> <li>Sterility: 0.22 µm filtered</li> <li>Endotoxin Level: Protein is endotoxin free.</li> <li>Grade: Crystallography grade</li> </ul>	Purification:	Two step purification of proteins expressed in baculovirus infected SF9 insect cells:
fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification si through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget Details		
through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.Purity:>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.Sterility:0.22 µm filteredEndotoxin Level:Protein is endotoxin free.Grade:Crystallography gradeTarget Details		
Sterility:     0.22 µm filtered       Endotoxin Level:     Protein is endotoxin free.       Grade:     Crystallography grade       Target Details     Vertical data data data data data data data da		
Endotoxin Level:     Protein is endotoxin free.       Grade:     Crystallography grade       Target Details     Value	Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Grade:     Crystallography grade       Target Details	Sterility:	0.22 µm filtered
Target Details	Endotoxin Level:	Protein is endotoxin free.
	Grade:	Crystallography grade
Target: TGFB2	Target Details	
	Target:	TGFB2

l arget:	IGFB2
Alternative Name:	TGFB2 (TGFB2 Products)
Background:	TGF-beta 2 has suppressive effects on interleukin-2 dependent T-cell growth.
Molecular Weight:	13.7 kDa Including tag.
UniProt:	P61812
Pathways:	Cell-Cell Junction Organization, Production of Molecular Mediator of Immune Response,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN3074159 | 02/07/2024 | Copyright antibodies-online. All rights reserved.

#### Protein targeting to Nucleus

## Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

### Images



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process