antibodies

Datasheet for ABIN3074869 TSSK1B Protein (AA 1-367) (Strep Tag)





Overview

Quantity:	1 mg
Target:	TSSK1B
Protein Characteristics:	AA 1-367
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This TSSK1B protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Sequence:	MDDAAVLKRR GYLLGINLGE GSYAKVKSAY SERLKFNVAI KIIDRKKAPA DFLEKFLPRE
	IEILAMLNHC SIIKTYEIFE TSHGKVYIVM ELAVQGDLLE LIKTRGALHE DEARKKFHQL
	SLAIKYCHDL DVVHRDLKCD NLLLDKDFNI KLSDFSFSKR CLRDDSGRMA LSKTFCGSPA
	YAAPEVLQGI PYQPKVYDIW SLGVILYIMV CGSMPYDDSN IKKMLRIQKE HRVNFPRSKH
	LTGECKDLIY HMLQPDVNRR LHIDEILSHC WMQPKARGSP SVAINKEGES SRGTEPLWTP
	EPGSDKKSAT KLEPEGEAQP QAQPETKPEG TAMQMSRQSE ILGFPSKPST METEEGPPQQ
	PPETRAQ
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN3074869 | 04/16/2024 | Copyright antibodies-online. All rights reserved.

- · Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALICE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3074869 | 04/16/2024 | Copyright antibodies-online. All rights reserved.

Purity:>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western BidEndotoxin Level:Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)Grade:Crystallography gradeTarget DetailsTSSK1BAlternative Name:TSSK1B (TSSK1B Products)Background:Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (Testis-specific serine/threonine-protein kinase 22A),FUNCTION: Testis-specific serine/threonine-protein kinase 22A),FUNCTION: Testis-specific serine/threonine-protein kinase 22A),FUNCTION: Testis-specific serine/threonine-protein kinase 22A),FUNCTION: Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (Testis-specific serine/threonine-protein kinase 22A),FUNCTION: Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (TON); Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (TON); Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (TON); Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TON); Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (TON); Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (TON); Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (TSK1) (TSSK-1) (TSK1) (TSSK-1) (TSK1) (TSSK-1) (TSK1) (TSSK-1) (TSSK1) (TSK1) (TSSK-1) (TSK1) (TSK1	
Grade: Crystallography grade Target Details TsSK1B Target: TSSK1B (TSSK1B Products) Alternative Name: TSSK1B (TSSK1B Products) Background: Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (Testis-speckinase 1) (EC 2.7.11.1) (Serine/threonine-protein kinase 22A),FUNCTION: Testis-speckinase 1) (EC 2.7.11.1) (Serine/threonine-protein kinase 22A),FUNCTION: Testis-speckinase 288' of TSKS. Involved in the late stages of spermatogenesis, during the reconstructic cytoplasm. During spermatogenesis, required for the transformation of a ring-shaped around the base of the flagellum originating from the chromatoid body. (ECO:0000269 PubMed:15733851, ECO:0000269 PubMed:19530700). Molecular Weight: 41.6 kDa	t.
Target Details Target: TSSK1B Alternative Name: TSSK1B (TSSK1B Products) Background: Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSSK-1) (Testis-specific serine/threonine-protein kinase 22A),FUNCTION: Testis-specific serine/threonine-protein kinase 22A),FUNCTION: Testis-specific serine/threonine-protein kinase required during spermatid development. Phosphoryla 288' of TSKS. Involved in the late stages of spermatogenesis, during the reconstructic cytoplasm. During spermatogenesis, required for the transformation of a ring-shaped around the base of the flagellum originating from the chromatoid body. (EC0:0000269 PubMed:15733851, EC0:0000269 PubMed:19530700). Molecular Weight: 41.6 kDa	
Target: TSSK1B Alternative Name: TSSK1B (TSSK1B Products) Background: Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (Testis-speckinase 1) (EC 2.7.11.1) (Serine/threonine-protein kinase 22A),FUNCTION: Testis-speckinase 1) (EC 2.7.11.1) (Serine/threonine-protein kinase 22A),FUNCTION: Testis-speckinase 288' of TSKS. Involved in the late stages of spermatid development. Phosphoryla 288' of TSKS. Involved in the late stages of spermatogenesis, during the reconstructic cytoplasm. During spermatogenesis, required for the transformation of a ring-shaped around the base of the flagellum originating from the chromatoid body. (ECO:0000269 PubMed:15733851, ECO:0000269 PubMed:19530700). Molecular Weight: 41.6 kDa	
Alternative Name: TSSK1B (TSSK1B Products) Background: Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSSK-1) (Testis-specific serine/threonine-protein kinase 22A),FUNCTION: Testis-specific serine/threonine-protein kinase required during spermatid development. Phosphoryla 288' of TSKS. Involved in the late stages of spermatogenesis, during the reconstructic cytoplasm. During spermatogenesis, required for the transformation of a ring-shaped around the base of the flagellum originating from the chromatoid body. (ECO:0000269 PubMed:15733851, ECO:0000269 PubMed:19530700}. Molecular Weight: 41.6 kDa	
Background: Testis-specific serine/threonine-protein kinase 1 (TSK-1) (TSK1) (TSSK-1) (Testis-specific serine/threonine-protein kinase 22A),FUNCTION: Testis-specific serine/threonine-protein kinase required during spermatid development. Phosphoryla 288' of TSKS. Involved in the late stages of spermatogenesis, during the reconstructic cytoplasm. During spermatogenesis, required for the transformation of a ring-shaped around the base of the flagellum originating from the chromatoid body. (ECO:0000269 PubMed:15733851, ECO:0000269 PubMed:19530700). Molecular Weight: 41.6 kDa	
kinase 1) (EC 2.7.11.1) (Serine/threonine-protein kinase 22A),FUNCTION: Testis-spect serine/threonine-protein kinase required during spermatid development. Phosphoryla 288' of TSKS. Involved in the late stages of spermatogenesis, during the reconstructi cytoplasm. During spermatogenesis, required for the transformation of a ring-shaped around the base of the flagellum originating from the chromatoid body. {ECO:0000269 PubMed:15733851, ECO:0000269 PubMed:19530700}. Molecular Weight: 41.6 kDa	
	ific ites 'Ser- on of the
UniProt: Q9BXA7	
Application Details	
Application Notes: In addition to the applications listed above we expect the protein to work for function as well. As the protein has not been tested for functional studies yet we cannot offer guarantee though.	
Comment: ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained Nicotiana tabacum c.v This contains all the protein expression machinery needed to even the most difficult-to-express proteins, including those that require post-translati modifications. During lysate production, the cell wall and other cellular components that are not req protein production are removed, leaving only the protein production machinery and th mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added t something that functions like a cell, but without the constraints of a living system - al needed is the DNA that codes for the desired protein!	o produce onal uired for ne o produce
Restrictions: For Research Use only	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN3074869 | 04/16/2024 | Copyright antibodies-online. All rights reserved.

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process