

Datasheet for ABIN3075497

VSX2 Protein (AA 1-361) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	VSX2
Protein Characteristics:	AA 1-361
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This VSX2 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Brand:	AlIcE®
Sequence:	<p>MTGKAGEALS KPKSETVAKS TSGGAPARCT GFGIQEILGL NKEPPSSHPR AALDGLAPGH LLAARSVLSP AGVGGMGLLG PGGLPGFYTQ PTFLEVLSDP QSVHLQPLGR ASGPLDTSQT ASSDSEDVSS SDRKMSKSAL NQTKKRKKRR HRTIFTSYQL EELEKAFNEA HYPDVYAREM LAMKTELPED RIQVWFQNRN AKWRKREKCW GRSSVMAEYG LYGAMVRHSI PLPESILKSA KDGIMDSCAP WLLGMHKKSL EAAAESGRKP EGERQALPKL DKMEQDERGP DAQAAISQEE LRENSIAVLR AKAQEHSTKV LGTVSGPDSL ARSTEKPEEE EAMDEDPAE RLSPQLEDMA</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>
Characteristics:	Key Benefits:

Product Details

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	VSX2
---------	------

Target Details

Alternative Name: VSX2 ([VSX2 Products](#))

Background: Visual system homeobox 2 (Ceh-10 homeodomain-containing homolog) (Homeobox protein CHX10),FUNCTION: Acts as a transcriptional regulator through binding to DNA at the consensus sequence 5'-[TC]TAATT[AG][AG]-3' upstream of gene promoters (PubMed:27301076). Plays a significant role in the specification and morphogenesis of the sensory retina (By similarity). May play a role in specification of V2a interneurons during spinal cord development (By similarity). Mediates differentiation of V2a interneurons by repression of motor neuron gene transcription, via competitively binding to response elements that are activated by the ISL1-LHX3 complex, such as VSX1 (PubMed:17919464, PubMed:27477290). Acts as a positive transcriptional regulator of NXNL1, regulation is significantly increased in synergy with VSX1 (By similarity). Acts as a negative transcriptional regulator of MITF (By similarity). Represses SAG transcription by competitive inhibition of ISL1-LHX3 response elements (PubMed:16236706, PubMed:27477290). Binds to the photoreceptor conserved element-1 (PCE-1) in the promoter of rod photoreceptor arrestin SAG and acts as a transcriptional repressor (By similarity). Involved in the development of retinal ganglion cells (RGCs) which leads to release of SHH by RGCs, promoting Hedgehog signaling and subsequent proliferation of retinal progenitor cells (By similarity). Participates in the development of the cells of the inner nuclear layer, by promoting postnatal differentiation of bipolar cells with a comparable inhibition of rod cell differentiation (By similarity). May play a role in the maintenance of neural retina identity during development by regulation of canonical Wnt genes and CTNNB1 localization, suggesting a role in the regulation of canonical Wnt signaling (PubMed:27301076). {ECO:0000250|UniProtKB:Q61412, ECO:0000269|PubMed:16236706, ECO:0000269|PubMed:17919464, ECO:0000269|PubMed:27301076, ECO:0000269|PubMed:27477290}.

Molecular Weight: 39.4 kDa

UniProt: [P58304](#)

Pathways: [Dopaminergic Neurogenesis](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from

Application Details

Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months