

Datasheet for ABIN3078088

CYP17A1 Protein (AA 1-508) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	CYP17A1
Protein Characteristics:	AA 1-508
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CYP17A1 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Brand:	AliCE®
Sequence:	<p>MWELVALLLL TLAYLFWPKR RCPGAKYPKS LLSLPLVGSL PFLPRHGHMH NNFFKLQKKY GPIYSVRMGT KTTVIVGHHQ LAKEVLIKKG KDFSGRPQMA TLDIASNNRK GIAFADSGAH WQLHRRRLAMA TFALFKDGDQ KLEKIICQEI STLCDMLATH NGQSIDISFP VFVAVTNVIS LICFNTSYKN GDPELNVIQN YNEGIIDNLS KDSLVDLVPW LKIFPNKTLE KLKSHVKIRN DLLNKILENY KEKFRSDSIT NMLDTLMQAK MNSDNGNAGP DQDSELLSDN HILTTIGDIF GAGVETTTSV VKWTLAFLH NPQVKKKLYE EIDQNVGFSR TPTISDRNRL LLEATIREV LRLRPVAPML IPHKANVDSS IGEFAVDKGT EVIINLWALH HNEKEWHQPD QFMPEFLNP AGTQLISPSV SYLPFGAGPR SCIGEILARQ ELFLIMAWLL QRFDLEVPDD GQLPSLEGIP KVVFLIDSKF VKIKVRQAWR EAQAEGST</p>

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade:

custom-made

Target Details

Target:	CYP17A1
Alternative Name:	CYP17A1 (CYP17A1 Products)
Background:	<p>Steroid 17-alpha-hydroxylase/17,20 lyase (EC 1.14.14.19) (17-alpha-hydroxyprogesterone aldolase) (EC 1.14.14.32) (CYPXVII) (Cytochrome P450 17A1) (Cytochrome P450-C17) (Cytochrome P450c17) (Steroid 17-alpha-monooxygenase),FUNCTION: A cytochrome P450 monooxygenase involved in corticoid and androgen biosynthesis (PubMed:9452426, PubMed:27339894, PubMed:22266943, PubMed:25301938). Catalyzes 17-alpha hydroxylation of C21 steroids, which is common for both pathways. A second oxidative step, required only for androgen synthesis, involves an acyl-carbon cleavage. The 17-alpha hydroxy intermediates, as part of adrenal glucocorticoids biosynthesis pathway, are precursors of cortisol (PubMed:9452426, PubMed:25301938) (Probable). Hydroxylates steroid hormones, pregnenolone and progesterone to form 17-alpha hydroxy metabolites, followed by the cleavage of the C17-C20 bond to form C19 steroids, dehydroepiandrosterone (DHEA) and androstenedione (PubMed:9452426, PubMed:27339894, PubMed:22266943, PubMed:25301938, PubMed:36640554). Has 16-alpha hydroxylase activity. Catalyzes 16-alpha hydroxylation of 17-alpha hydroxy pregnenolone, followed by the cleavage of the C17-C20 bond to form 16-alpha-hydroxy DHEA (PubMed:36640554). Also 16-alpha hydroxylates androgens, relevant for estriol synthesis (PubMed:27339894, PubMed:25301938). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR, NADPH-ferrihemoprotein reductase) (PubMed:9452426, PubMed:27339894, PubMed:22266943, PubMed:25301938). {ECO:0000269 PubMed:22266943, ECO:0000269 PubMed:25301938, ECO:0000269 PubMed:27339894, ECO:0000269 PubMed:36640554, ECO:0000269 PubMed:9452426, ECO:0000305 PubMed:8027220}.</p>
Molecular Weight:	57.4 kDa
UniProt:	P05093
Pathways:	Metabolism of Steroid Hormones and Vitamin D , Steroid Hormone Biosynthesis , Regulation of Hormone Metabolic Process , Regulation of Hormone Biosynthetic Process , C21-Steroid Hormone Metabolic Process , Cellular Response to Molecule of Bacterial Origin

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
--------------------	---

Application Details

as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment:

ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:

Liquid

Buffer:

The buffer composition is at the discretion of the manufacturer.

Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**

Handling Advice:

Avoid repeated freeze-thaw cycles.

Storage:

-80 °C

Storage Comment:

Store at -80°C.

Expiry Date:

12 months