

Datasheet for ABIN3078308

CYP3A7 Protein (AA 1-503) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	CYP3A7
Protein Characteristics:	AA 1-503
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CYP3A7 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Brand:	AliCE®
Sequence:	<p>MDLIPNLAVE TWLLAVSLI LLYLYGTRTH GLFKKLGIPG PTPLPFLGNA LSFRKGYWTF DMECYKKYRK VWGIYDCQP MLAITDPDMI KTVLVKECYS VFTNRRPFGP VGFMKNAISI AEDEEWKRIR SLLSPTFTSG KLKEMVPIIA QYGDVLRNL RREAETGKPV TLKHVFGAYS MDVITSTSFG VSIDSLNNPQ DPFVENTKKL LRFNPLDPFV LSIKVPFLT PILEALNITV FPRKVISFLT KSVKQIKEGR LKETQKHRVD FLQLMIDSQN SKDSETHKAL SDLELMAQSI IFIFAGYETT SSVLSFIIYE LATHPDVQQK VQKEIDTVLP NKAPPTYDTV LQLEYLDMVV NETLRLFPVA MRLERVCKKD VEINGMFIPK GVVMIPSYV LHHPKPYWTE PEKFLPERFS KKNKDNIDPY IYTPFGSGPR NCIGMRFALV NMKLALVRVL QNFSFKPCKE TQIPLKLRFG GLLLTEKPIV LKAESRDETV SGA</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you</p>

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade:

custom-made

Target Details

Target:	CYP3A7
Alternative Name:	CYP3A7 (CYP3A7 Products)
Background:	<p>Cytochrome P450 3A7 (EC 1.14.14.1) (CYP11A7) (Cytochrome P450-HFLA) (P450HLP2),FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of steroid hormones and vitamins during embryogenesis (PubMed:9555064, PubMed:11093772, PubMed:14559847, PubMed:12865317, PubMed:17178770). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:9555064, PubMed:11093772, PubMed:14559847, PubMed:12865317, PubMed:17178770). Catalyzes the hydroxylation of carbon-hydrogen bonds. Metabolizes 3beta-hydroxyandrost-5-en-17-one (dehydroepiandrosterone, DHEA), a precursor in the biosynthesis of androgen and estrogen steroid hormones (PubMed:9555064, PubMed:17178770). Exhibits high catalytic activity for the formation of hydroxyestrogens from estrone (E1), particularly D-ring hydroxylated estrone at the C16-alpha position (PubMed:14559847, PubMed:12865317). Mainly hydroxylates all trans-retinoic acid (atRA) to 4-hydroxyretinoate and may play a role in atRA clearance during fetal development (PubMed:11093772). Also involved in the oxidative metabolism of xenobiotics including anticonvulsants (PubMed:9555064). {ECO:0000269 PubMed:11093772, ECO:0000269 PubMed:12865317, ECO:0000269 PubMed:14559847, ECO:0000269 PubMed:17178770, ECO:0000269 PubMed:9555064}.</p>
Molecular Weight:	57.5 kDa
UniProt:	P24462
Pathways:	Steroid Hormone Biosynthesis

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	<p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for</p>

Application Details

protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months