
# antibodies .- online.com





# CYP3A7 Protein (AA 1-503) (Strep Tag)



### Overview

Quantity:	1 mg
Target:	CYP3A7
Protein Characteristics:	AA 1-503
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CYP3A7 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

### **Product Details**

Sequence:

MDLIPNLAVE TWLLLAVSLI LLYLYGTRTH GLFKKLGIPG PTPLPFLGNA LSFRKGYWTF
DMECYKKYRK VWGIYDCQQP MLAITDPDMI KTVLVKECYS VFTNRRPFGP VGFMKNAISI
AEDEEWKRIR SLLSPTFTSG KLKEMVPIIA QYGDVLVRNL RREAETGKPV TLKHVFGAYS
MDVITSTSFG VSIDSLNNPQ DPFVENTKKL LRFNPLDPFV LSIKVFPFLT PILEALNITV
FPRKVISFLT KSVKQIKEGR LKETQKHRVD FLQLMIDSQN SKDSETHKAL SDLELMAQSI
IFIFAGYETT SSVLSFIIYE LATHPDVQQK VQKEIDTVLP NKAPPTYDTV LQLEYLDMVV
NETLRLFPVA MRLERVCKKD VEINGMFIPK GVVVMIPSYV LHHDPKYWTE PEKFLPERFS
KKNKDNIDPY IYTPFGSGPR NCIGMRFALV NMKLALVRVL QNFSFKPCKE TQIPLKLRFG
GLLLTEKPIV LKAESRDETV SGA

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and

### **Product Details**

	Western blot.
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Target Details	
Target:	CYP3A7
Alternative Name:	CYP3A7 (CYP3A7 Products)
Background:	Cytochrome P450 3A7 (EC 1.14.14.1) (CYPIIIA7) (Cytochrome P450-HFLA)
	(P450HLp2),FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of
	steroid hormones and vitamins during embryogenesis (PubMed:9555064, PubMed:11093772,
	PubMed:14559847, PubMed:12865317, PubMed:17178770). Mechanistically, uses molecular
	oxygen inserting one oxygen atom into a substrate, and reducing the second into a water
	molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH
	hemoprotein reductase) (PubMed:9555064, PubMed:11093772, PubMed:14559847,
	PubMed:12865317, PubMed:17178770). Catalyzes the hydroxylation of carbon-hydrogen
	bonds. Metabolizes 3beta-hydroxyandrost-5-en-17-one (dehydroepiandrosterone, DHEA), a
	precursor in the biosynthesis of androgen and estrogen steroid hormones (PubMed:9555064,
	PubMed:17178770). Exhibits high catalytic activity for the formation of hydroxyestrogens from
	estrone (E1), particularly D-ring hydroxylated estrone at the C16-alpha position
	(PubMed:14559847, PubMed:12865317). Mainly hydroxylates all trans-retinoic acid (atRA) to 4
	hydroxyretinoate and may play a role in atRA clearance during fetal development
	(PubMed:11093772). Also involved in the oxidative metabolism of xenobiotics including
	anticonvulsants (PubMed:9555064). {ECO:0000269 PubMed:11093772,
	ECO:0000269 PubMed:12865317, ECO:0000269 PubMed:14559847,
	ECO:0000269 PubMed:17178770, ECO:0000269 PubMed:9555064}.
Molecular Weight:	57.5 kDa
JniProt:	P24462
Pathways:	Steroid Hormone Biosynthesis
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a

## **Application Details**

Application betails		
	guarantee though.	
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.  During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!	
Restrictions: Handling	For Research Use only	
Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	Unlimited (if stored properly)	