
antibodies .- online.com





CYP7A1 Protein (AA 1-504) (Strep Tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	CYP7A1
Protein Characteristics:	AA 1-504
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CYP7A1 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Sequence:

MMTTSLIWGI AIAACCCLWL ILGIRRRQTG EPPLENGLIP YLGCALQFGA NPLEFLRANQ RKHGHVFTCK LMGKYVHFIT NPLSYHKVLC HGKYFDWKKF HFATSAKAFG HRSIDPMDGN TTENINDTFI KTLQGHALNS LTESMMENLQ RIMRPPVSSN SKTAAWVTEG MYSFCYRVMF EAGYLTIFGR DLTRRDTQKA HILNNLDNFK QFDKVFPALV AGLPIHMFRT AHNAREKLAE SLRHENLQKR ESISELISLR MFLNDTLSTF DDLEKAKTHL VVLWASQANT IPATFWSLFQ MIRNPEAMKA ATEEVKRTLE NAGQKVSLEG NPICLSQAEL NDLPVLDSII KESLRLSSAS LNIRTAKEDF TLHLEDGSYN IRKDDIIALY PQLMHLDPEI YPDPLTFKYD RYLDENGKTK TTFYCNGLKL KYYYMPFGSG ATICPGRLFA IHEIKQFLIL MLSYFELELI EGQAKCPPLD QSRAGLGILP PLNDIEFKYK FKHL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and

Product Details

	Western blot.
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade
Target Details	
Target:	CYP7A1
Alternative Name:	CYP7A1 (CYP7A1 Products)
Background:	Cytochrome P450 7A1 (24-hydroxycholesterol 7-alpha-hydroxylase) (EC 1.14.14.26) (CYPVII)
	(Cholesterol 7-alpha-hydroxylase) (Cholesterol 7-alpha-monooxygenase) (EC
	1.14.14.23),FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of
	endogenous cholesterol and its oxygenated derivatives (oxysterols) (PubMed:11013305,
	PubMed:12077124, PubMed:19965590, PubMed:2384150, PubMed:21813643).
	Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and
	reducing the second into a water molecule, with two electrons provided by NADPH via
	cytochrome P450 reductase (CPR, NADPH-ferrihemoprotein reductase) (PubMed:2384150,
	PubMed:11013305, PubMed:12077124, PubMed:19965590, PubMed:21813643). Functions as
	a critical regulatory enzyme of bile acid biosynthesis and cholesterol homeostasis. Catalyzes
	the hydroxylation of carbon hydrogen bond at 7-alpha position of cholesterol, a rate-limiting
	step in cholesterol catabolism and bile acid biosynthesis (PubMed:12077124,
	PubMed:19965590, PubMed:2384150). 7-alpha hydroxylates several oxysterols, including
	4beta-hydroxycholesterol and 24-hydroxycholesterol (PubMed:11013305, PubMed:12077124)
	Catalyzes the oxidation of the 7,8 double bond of 7-dehydrocholesterol and lathosterol with
	direct and predominant formation of the 7-keto derivatives (PubMed:21813643).
	{ECO:0000269 PubMed:11013305, ECO:0000269 PubMed:12077124,
	ECO:0000269 PubMed:19965590, ECO:0000269 PubMed:21813643,
	ECO:0000269 PubMed:2384150}.
Molecular Weight:	57.7 kDa
UniProt:	P22680
Pathways:	Steroid Hormone Biosynthesis, Carbohydrate Homeostasis, Regulation of Lipid Metabolism by
	PPARalpha

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for
	protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process