antibodies -online.com





COPS7A Protein (AA 2-275) (His tag)



Image



Go to Product page

()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	1 mg
Target:	COPS7A
Protein Characteristics:	AA 2-275
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This COPS7A protein is labelled with His tag.
Application:	SDS-PAGE (SDS), ELISA, Western Blotting (WB), Crystallization (Crys)
Product Details	

Purification tag / Conjugate:	This COPS7A protein is labelled with His tag.	
Application:	SDS-PAGE (SDS), ELISA, Western Blotting (WB), Crystallization (Crys)	
Product Details		
Sequence:	SAEVKVTGQN QEQFLLLAKS AKGAALATLI HQVLEAPGVY VFGELLDMPN VRELAESDFA	
	STFRLLTVFA YGTYADYLAE ARNLPPLTEA QKNKLRHLSV VTLAAKVKCI PYAVLLEALA	
	LRNVRQLEDL VIEAVYADVL RGSLDQRNQR LEVDYSIGRD IQRQDLSAIA RTLQEWCVGC	
	EVVLSGIEEQ VSRANQHKEQ QLGLKQQIES EVANLKKTIK VTTAAAAAAT SQDPEQHLTE	
	LREPAPGTNQ RQPSKKASKG KGLRGSAKIW SKSN	
	Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a	
	special request, please contact us.	
Characteristics:	Made in Germany - from design to production - by highly experienced protein experts.	
	 Human COPS7A Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. 	
	State-of-the-art algorithm used for plasmid design (Gene synthesis).	
	This protein is a made to order protein and will be made for the first time for your order. Our	

experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

 $0.22\,\mu m$ filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

Target Details

Target:	COPS7A
Alternative Name:	COPS7A (COPS7A Products)
Background:	Component of the COP9 signalosome complex (CSN), a complex involved in various cellular
	and developmental processes. The CSN complex is an essential regulator of the ubiquitin (UbI)

	conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3
	ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as
	SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, JUN, I-kappa-
	B-alpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases.
	CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the
	Ubl system, respectively. {ECO:0000269 PubMed:11285227, ECO:0000269 PubMed:11337588,
	ECO:0000269 PubMed:12628923, ECO:0000269 PubMed:12732143,
	ECO:0000269 PubMed:9535219}.
Molecular Weight:	31.1 kDa Including tag.
UniProt:	Q9UBW8

Cell Division Cycle

Application Details

Pathways:

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be
	insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to
	increase solubility. We will discuss all possible options with you in detail to assure that you
	receive your protein of interest.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process