antibodies - online.com







CXCL7 Protein (AA 59-121) (His tag)



Image



\sim				
	11/6	٦r١	/10	۱۸.

1 mg	
CXCL7 (PPBP)	
AA 59-121	
Human	
Escherichia coli (E. coli)	
Recombinant	
This CXCL7 protein is labelled with His tag.	
Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)	
AELRCMCIKT TSGIHPKNIQ SLEVIGKGTH CNQVEVIATL KDGRKICLDP DAPRIKKIVQ KKL	
Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a	
special request, please contact us.	
 Made in Germany - from design to production - by highly experienced protein experts. Human PPBP Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis). 	
This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.	

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

protein experts will do everything to make sure that you receive the protein you ordered.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

CXCL7 (PPBP)

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Target:

Crystallography grade

Target Details

Alternative Name:	PPBP (PPBP Products)
Background:	LA-PF4 stimulates DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation,
	prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It
	also stimulates the formation and secretion of plasminogen activator by human synovial cells.
	NAP-2 is a ligand for CXCR1 and CXCR2, and NAP-2, NAP-2(73), NAP-2(74), NAP-2(1-66), and
	most potent NAP-2(1-63) are chemoattractants and activators for neutrophils. TC-1 and TC-2
	are antibacterial proteins, in vitro released from activated platelet alpha-granules. CTAP-III(1-81)

Target Details

Storage Comment:

Expiry Date:

l arget Details		
	is more potent than CTAP-III desensitize chemokine-induced neutrophil activation. {ECO:0000269 PubMed:10877842, ECO:0000269 PubMed:7890771, ECO:0000269 PubMed:8950790, ECO:0000269 PubMed:9794434}.	
Molecular Weight:	7.9 kDa Including tag.	
UniProt:	P02775	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.	
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	

Store at -80°C.

Unlimited (if stored properly)



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process