

Datasheet for ABIN3078713

DAZAP2 Protein (AA 1-168) (Strep Tag)[Go to Product page](#)**1** Image

Overview

| | |
|-------------------------------|---|
| Quantity: | 1 mg |
| Target: | DAZAP2 |
| Protein Characteristics: | AA 1-168 |
| Origin: | Human |
| Source: | Tobacco (Nicotiana tabacum) |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This DAZAP2 protein is labelled with Strep Tag. |
| Application: | ELISA, Western Blotting (WB), SDS-PAGE (SDS) |

Product Details

| | |
|------------------|--|
| Sequence: | <p>MNSKGQYPTQ PTYPVQPPGN PVYPQTLHLP QAPPYTDAPP AYSELYRPSF VHPGAATVPT MSAAFPGLSL YLPMAQSVAV GPLGSTIPMA YYPVGPIYPP GSTVLVEGGY DAGARFGAGA TAGNIPPPPP GCPPNAAQLA VMQGANVLVT QRKGNFFMGG SDGGYTIW</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p> |
| Characteristics: | <p>Key Benefits:</p> <ul style="list-style-type: none">• Made in Germany - from design to production - by highly experienced protein experts.• Protein expressed with ALICE® and purified by multi-step, protein-specific process to ensure correct folding and modification.• These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed). |

- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

| | |
|------------------|--|
| Purification: | Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®): 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot. |
| Purity: | >80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. |
| Endotoxin Level: | Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg) |
| Grade: | Crystallography grade |

Target Details

| | |
|-------------------|--|
| Target: | DAZAP2 |
| Alternative Name: | DAZAP2 (DAZAP2 Products) |
| Background: | <p>DAZ-associated protein 2 (Deleted in azoospermia-associated protein 2) (Proline-rich transcript in brain protein),FUNCTION: In unstressed cells, promotes SIAH1-mediated polyubiquitination and degradation of the serine/threonine-protein kinase HIPK2, probably by acting as a loading factor that potentiates complex formation between HIPK2 and ubiquitin ligase SIAH1 (PubMed:33591310). In response to DNA damage, localizes to the nucleus following phosphorylation by HIPK2 and modulates the expression of a subset of TP53/p53 target genes by binding to TP53 at target gene promoters (PubMed:33591310). This limits the expression of a number of cell death-mediating TP53 target genes, reducing DNA damage-induced cell death (PubMed:33591310). Enhances the binding of transcription factor TCF7L2/TCF4, a Wnt signaling pathway effector, to the promoters of target genes (By similarity). Plays a role in stress granule formation (PubMed:17984221). {ECO:0000250 UniProtKB:Q9DCP9, ECO:0000269 PubMed:17984221, ECO:0000269 PubMed:33591310}.</p> |
| Molecular Weight: | 17.3 kDa |
| UniProt: | Q15038 |

Application Details

| | |
|--------------------|--|
| Application Notes: | <p>In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.</p> |
| Comment: | <p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!</p> |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|--|
| Format: | Liquid |
| Buffer: | The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. |
| Expiry Date: | Unlimited (if stored properly) |

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process