

Datasheet for ABIN3078974

APPL2 Protein (AA 1-664) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	APPL2
Protein Characteristics:	AA 1-664
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This APPL2 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Brand:	ALICE®
Sequence:	<p>MPAVDKLLLE EALQDSPQTR SLLSVFEEDA GTLTDYTNQL LQAMQRVYGA QNEMCLATQQ</p> <p>LSKQLLAYEK QNFALGKGDE EVISTLHYFS KVVDELNLLH TELAKQLADT MVLPIIQFRE</p> <p>KDLTEVSTLK DLFGLASNEH DLSMAKYSRL PKKKENEKVK TEVGKEVAAA RRKQHLSSLQ</p> <p>YYCALNALQY RKQMAMMEPM IGFAHGQINF FKKGAEMFSK RMDSFLSSVA DMVQSIQVEL</p> <p>EAEAEKMRVS QQELLSVDES VYTPDSDVAA PQINRNLIQK AGYLNLRNKT GLVTTTWERL</p> <p>YFFTQGGNLM CQPRGAVAGG LIQDLNCSV MAVDCEDRRY CFQITTPNGK SGILQAESR</p> <p>KENEEWICAI NNISRQIYLT DNPEAVAIKL NQTALQAVTP ITSFGKKQES SCPSQNLKNS</p> <p>EMENENDKIV PKATASLPEA EELIAPGTPI QFDIVLPATE FLDQNRGSRR TNPFGETEDE</p> <p>SFPEAEDSLL QQMFIIVRFLG SMAVKTDSTT EVIYEAMRQV LAARAIHNIF RMTESHLMVT</p> <p>SQSLRLIDPQ TQVSRANFEL TSVTQFAAHQ ENKRLVGFVI RVPESTGEES LSTYIFESNS</p> <p>EGEKICYAIN LGKEIIEVQK DPEALAQLML SIPLTNDGKY VLLNDQPDDD DGPNNEHRGA ESEA</p>

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Product Details

Grade: custom-made

Target Details

Target: APPL2

Alternative Name: APPL2 ([APPL2 Products](#))

Background: DCC-interacting protein 13-beta (Dip13-beta) (Adapter protein containing PH domain, PTB domain and leucine zipper motif 2),FUNCTION: Multifunctional adapter protein that binds to various membrane receptors, nuclear factors and signaling proteins to regulate many processes, such as cell proliferation, immune response, endosomal trafficking and cell metabolism (PubMed:26583432, PubMed:15016378, PubMed:24879834). Regulates signaling pathway leading to cell proliferation through interaction with RAB5A and subunits of the NuRD/MeCP1 complex (PubMed:15016378). Plays a role in immune response by modulating phagocytosis, inflammatory and innate immune responses. In macrophages, enhances Fc-gamma receptor-mediated phagocytosis through interaction with RAB31 leading to activation of PI3K/Akt signaling. In response to LPS, modulates inflammatory responses by playing a key role on the regulation of TLR4 signaling and in the nuclear translocation of RELA/NF-kappa-B p65 and the secretion of pro- and anti-inflammatory cytokines. Also functions as a negative regulator of innate immune response via inhibition of AKT1 signaling pathway by forming a complex with APPL1 and PIK3R1 (By similarity). Plays a role in endosomal trafficking of TGFBR1 from the endosomes to the nucleus (PubMed:26583432). Plays a role in cell metabolism by regulating adiponectin and insulin signaling pathways and adaptive thermogenesis (PubMed:24879834) (By similarity). In muscle, negatively regulates adiponectin-stimulated glucose uptake and fatty acid oxidation by inhibiting adiponectin signaling pathway through APPL1 sequestration thereby antagonizing APPL1 action (By similarity). In muscles, negatively regulates insulin-induced plasma membrane recruitment of GLUT4 and glucose uptake through interaction with TBC1D1 (PubMed:24879834). Plays a role in cold and diet-induced adaptive thermogenesis by activating ventromedial hypothalamus (VMH) neurons through AMPK inhibition which enhances sympathetic outflow to subcutaneous white adipose tissue (sWAT), sWAT browning and cold tolerance (By similarity). Also plays a role in other signaling pathways namely Wnt/beta-catenin, HGF and glucocorticoid receptor signaling (PubMed:19433865) (By similarity). Positive regulator of beta-catenin/TCF-dependent transcription through direct interaction with RUVBL2/reptin resulting in the relief of RUVBL2-mediated repression of beta-catenin/TCF target genes by modulating the interactions within the beta-catenin-reptin-HDAC complex (PubMed:19433865). May affect adult neurogenesis in hippocampus and olfactory system via regulating the sensitivity of glucocorticoid receptor.

Target Details

Required for fibroblast migration through HGF cell signaling (By similarity).
{ECO:0000250|UniProtKB:Q8K3G9, ECO:0000269|PubMed:15016378,
ECO:0000269|PubMed:19433865, ECO:0000269|PubMed:24879834,
ECO:0000269|PubMed:26583432}.

Molecular Weight: 74.5 kDa

UniProt: [Q8NEU8](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.
Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol **Might differ depending on protein.**

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months