
antibodies .- online.com





DOM3Z Protein (AA 1-396) (Strep Tag)



Overview

Quantity:	1 mg	
Target:	DOM3Z	
Protein Characteristics:	AA 1-396	
Origin:	Human	
Source:	Tobacco (Nicotiana tabacum)	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This DOM3Z protein is labelled with Strep Tag.	
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)	

Product Details

Sequence:

MDPRGTKRGA EKTEVAEPRN KLPRPAPSLP TDPALYSGPF PFYRRPSELG CFSLDAQRQY
HGDARALRYY SPPPTNGPGP NFDLRDGYPD RYQPRDEEVQ ERLDHLLCWL LEHRGRLEGG
PGWLAEAIVT WRGHLTKLLT TPYERQEGWQ LAASRFQGTL YLSEVETPNA RAQRLARPPL
LRELMYMGYK FEQYMCADKP GSSPDPSGEV NTNVAFCSVL RSRLGSHPLL FSGEVDCTDP
QAPSTQPPTC YVELKTSKEM HSPGQWRSFY RHKLLKWWAQ SFLPGVPNVV AGFRNPDGFV
SSLKTFPTMK MFEYVRNDRD GWNPSVCMNF CAAFLSFAQS TVVQDDPRLV HLFSWEPGGP
VTVSVHQDAP YAFLPIWYVE AMTQDLPSPP KTPSPK

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you

have a special request, please contact us.

Characteristics: Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- · State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Product Details

Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.	
Endotoxin Level:	toxin Level: Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)	

Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)		
Target Details			
Target:	DOM3Z		
Alternative Name:	DXO (DOM3Z Products)		
Background:	Decapping and exoribonuclease protein (DXO) (EC 3.6.1) (5'-3' exoribonuclease DXO) (EC		
	3.1.13) (Dom-3 homolog Z) (NAD-capped RNA hydrolase DXO) (DeNADding enzyme DXO) (EC		
	3.6.1),FUNCTION: Decapping enzyme for NAD-capped RNAs: specifically hydrolyzes the		
	nicotinamide adenine dinucleotide (NAD) cap from a subset of RNAs by removing the entire		
	NAD moiety from the 5'-end of an NAD-capped RNA (PubMed:28283058). The NAD-cap is		
	present at the 5'-end of some RNAs and snoRNAs (PubMed:28283058). In contrast to the		
	canonical 5'-end N7 methylguanosine (m7G) cap, the NAD cap promotes mRNA decay		
	(PubMed:28283058). Preferentially acts on NAD-capped transcripts in response to		
	environmental stress (PubMed:31101919). Also acts as a non-canonical decapping enzyme		
	that removes the entire cap structure of m7G capped or incompletely capped RNAs and		
	mediates their subsequent degradation (By similarity). Specifically degrades pre-mRNAs with a		
	defective 5'-end m7G cap and is part of a pre-mRNA capping quality control (By similarity). Has		
	decapping activity toward incomplete 5'-end m7G cap mRNAs such as unmethylated 5'-end-		
	capped RNA (cap0), while it has no activity toward 2'-O-ribose methylated m7G cap (cap1)		
	(PubMed:29601584). In contrast to canonical decapping enzymes DCP2 and NUDT16, which		
	cleave the cap within the triphosphate linkage, the decapping activity releases the entire cap		
	structure GpppN and a 5'-end monophosphate RNA (By similarity). Also has 5'-3'		
	exoribonuclease activities: The 5'-end monophosphate RNA is then degraded by the 5'-3'		
	exoribonuclease activity, enabling this enzyme to decap and degrade incompletely capped		
	mRNAs (PubMed:29601584). Also possesses RNA 5'-pyrophosphohydrolase activity by		
	hydrolyzing the 5'-end triphosphate to release pyrophosphates (By similarity). Exhibits		
	decapping activity towards FAD-capped RNAs (PubMed:32374864). Exhibits decapping activity		
	towards dpCoA-capped RNAs in vitro (By similarity). {ECO:0000250 UniProtKB:070348,		
	ECO:0000269 PubMed:28283058, ECO:0000269 PubMed:29601584,		
	ECO:0000269 PubMed:31101919, ECO:0000269 PubMed:32374864}.		
Molecular Weight:	44.9 kDa		

Molecular Weight:	44.9 kDa
UniProt:	077932

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.	
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for	
	protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	Unlimited (if stored properly)	