

Datasheet for ABIN3082495

JMJD5 Protein (AA 1-416) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	JMJD5
Protein Characteristics:	AA 1-416
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This JMJD5 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Brand:	ALICE®
Sequence:	<p>MAGDTHCPAE PLAREGTLWE ALRALLPHSK EDLKLDLGEK VERSVVTLLQ RATELFYEGR RDECLQSSEV ILDYSWEKLN TGTWQDVDDK WRRVYAIGCL LKALCLCQAP EDANTVAAAL RVCDMGLLMG AAILGDILLK VAAILQTHLP GKRPARGLSLP EQPCTKKARA DHGLIPDVKL EKTVPRLHRP SLQHFREQFL VPGRPVILKG VADHWPCMOK WSLEYIQEIA GCRTVPVEVG SRYTDEEWSQ TLMTVNEFIS KYIVNEPRDV GYLAQHQLFD QIPELKQDIS IPDYCSLGDG EEEEITINAW FGPQGTISPL HQDPQQNFLV QVMGRKYIRL YSPQESGALY PHDTHLLHNT SQVDVENPDL EKFPKFAKAP FLSCILSPGE ILFIPVKYWH YVRALDLSFS VSFWWS</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>
Characteristics:	Key Benefits:

Product Details

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	JMJD5
---------	-------

Target Details

Alternative Name: KDM8 ([JMJD5 Products](#))

Background: Bifunctional peptidase and arginyl-hydroxylase JMJD5 (EC 1.14.11.73) (EC 3.4.-.-) (JmjC domain-containing protein 5) (Jumonji C domain-containing protein 5) (L-arginine (3R)-hydroxylase KDM8), FUNCTION: Bifunctional enzyme that acts both as an endopeptidase and 2-oxoglutarate-dependent monooxygenase (PubMed:28847961, PubMed:29459673, PubMed:28982940, PubMed:29563586). Endopeptidase that cleaves histones N-terminal tails at the carboxyl side of methylated arginine or lysine residues, to generate 'tailless nucleosomes', which may trigger transcription elongation (PubMed:28847961, PubMed:29459673, PubMed:28982940). Preferentially recognizes and cleaves monomethylated and dimethylated arginine residues of histones H2, H3 and H4. After initial cleavage, continues to digest histones tails via its aminopeptidase activity (PubMed:28847961, PubMed:29459673). Upon DNA damage, cleaves the N-terminal tail of histone H3 at monomethylated lysine residues, preferably at monomethylated 'Lys-9' (H3K9me1). The histone variant H3F3A is the major target for cleavage (PubMed:28982940). Additionally, acts as a Fe(2+) and 2-oxoglutarate-dependent monooxygenase, catalyzing (R)-stereospecific hydroxylation at C-3 of 'Arg-137' of RPS6 and 'Arg-141' of RCCD1, but the biological significance of this activity remains to be established (PubMed:29563586). Regulates mitosis through different mechanisms: Plays a role in transcriptional repression of satellite repeats, possibly by regulating H3K36 methylation levels in centromeric regions together with RCCD1. Possibly together with RCCD1, is involved in proper mitotic spindle organization and chromosome segregation (PubMed:24981860). Negatively regulates cell cycle repressor CDKN1A/p21, which controls G1/S phase transition (PubMed:24740926). Required for G2/M phase cell cycle progression. Regulates expression of CCNA1/cyclin-A1, leading to cancer cell proliferation (PubMed:20457893). Also, plays a role in regulating alpha-tubulin acetylation and cytoskeletal microtubule stability involved in epithelial to mesenchymal transition (PubMed:28455245). Regulates the circadian gene expression in the liver (By similarity). Represses the transcriptional activator activity of the CLOCK-BMAL1 heterodimer in a catalytically-independent manner (PubMed:30500822). Negatively regulates the protein stability and function of CRY1, required for AMPK-FBXL3-induced CRY1 degradation (PubMed:30500822). {ECO:0000250|UniProtKB:Q9CXT6, ECO:0000269|PubMed:20457893, ECO:0000269|PubMed:24740926, ECO:0000269|PubMed:24981860, ECO:0000269|PubMed:28455245, ECO:0000269|PubMed:28847961, ECO:0000269|PubMed:28982940, ECO:0000269|PubMed:29459673, ECO:0000269|PubMed:29563586, ECO:0000269|PubMed:30500822}.

Molecular Weight: 47.3 kDa

Target Details

UniProt:	Q8N371
Pathways:	Chromatin Binding

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	<p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	<p>The buffer composition is at the discretion of the manufacturer.</p> <p>Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.</p>
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months