

Datasheet for ABIN308258
anti-DAPP1 antibody (C-Term)[Go to Product page](#)

1 Image

Overview

| | |
|----------------------|--------------------------------------|
| Quantity: | 100 µg |
| Target: | DAPP1 |
| Binding Specificity: | C-Term |
| Reactivity: | Human |
| Host: | Goat |
| Clonality: | Polyclonal |
| Conjugate: | This DAPP1 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA |

Product Details

| | |
|-------------------|--|
| Purpose: | DAPP1 |
| Immunogen: | Peptide with sequence SALCISPEEKTDHK, from the C Terminus of the protein sequence according to AAF14578.1. |
| Sequence: | SALCISPEEK TDHK |
| Isotype: | IgG |
| Specificity: | This antibody was raised against the C-terminus of NP_055210.1. However, NCBI has changed the version to NP_055210.2 thus losing the C-terminal epitope. |
| Cross-Reactivity: | Human |
| Purification: | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |

Product Details

Grade: Verified

Target Details

Target: DAPP1

Alternative Name: DAPP1 ([DAPP1 Products](#))

Background: DAPP1, dual adaptor of phosphotyrosine and 3-phosphoinositides, BAM32, DKFZp667E0716, HSPC066, B-cell adapter molecule of 32 kDa, b lymphocyte adapter protein Bam32, dual adapter for phosphotyrosine and 3-phosphotyrosine and 3-phosphoinositide, hDAPP1

Gene ID: 27071

Pathways: [BCR Signaling](#)

Application Details

Application Notes: Western Blot: Approx 30 kDa band observed in Human Kidney lysates (calculated MW of 32.0 kDa according to AAF14578.1). Recommended concentration: 0.5-1 µg/mL.
Peptide ELISA: antibody detection limit dilution 1:4000.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 0.5 mg/mL

Buffer: Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Minimize freezing and thawing.

Storage: -20 °C

Storage Comment: Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Western Blotting

Image 1. ABIN308258 (0.5µg/ml) staining of Human Kidney lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.