antibodies

Datasheet for ABIN3083158 HBXIP Protein (AA 1-91) (Strep Tag)



Overview

Overview	
Quantity:	1 mg
Target:	HBXIP
Protein Characteristics:	AA 1-91
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This HBXIP protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)
Product Details	
Sequence:	MEATLEQHLE DTMKNPSIVG VLCTDSQGLN LGCRGTLSDE HAGVISVLAQ QAAKLTSDPT
	DIPVVCLESD NGNIMIQKHD GITVAVHKMA S
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:
	 Made in Germany - from design to production - by highly experienced protein experts. Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
	These proteins are normally active (enzymatically functional) as our customers have
	reported (not tested by us and not guaranteed).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):
	1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
	 Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Target Details	
Target:	HBXIP

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN3083158 | 05/07/2024 | Copyright antibodies-online. All rights reserved.

Alternative Name:	LAMTOR5 (HBXIP Products)
Target Type:	Viral Protein
Background:	
	Ragulator complex protein LAMTOR5 (Hepatitis B virus X-interacting protein) (HBV X-
	interacting protein) (HBX-interacting protein) (Late endosomal/lysosomal adaptor and MAPK
	and MTOR activator 5), FUNCTION: As part of the Ragulator complex it is involved in amino acid
	sensing and activation of mTORC1, a signaling complex promoting cell growth in response to
	growth factors, energy levels, and amino acids (PubMed:22980980, PubMed:30181260,
	PubMed:29158492). Activated by amino acids through a mechanism involving the lysosomal V
	ATPase, the Ragulator plays a dual role for the small GTPases Rag (RagA/RRAGA,
	RagB/RRAGB, RagC/RRAGC and/or RagD/RRAGD): it (1) acts as a guanine nucleotide
	exchange factor (GEF), activating the small GTPases Rag and (2) mediates recruitment of Rag
	GTPases to the lysosome membrane (PubMed:22980980, PubMed:30181260,
	PubMed:29107538, PubMed:29158492, PubMed:28935770). Activated Ragulator and Rag
	GTPases function as a scaffold recruiting mTORC1 to lysosomes where it is in turn activated
	(PubMed:22980980, PubMed:30181260, PubMed:29158492). When complexed to BIRC5,
	interferes with apoptosome assembly, preventing recruitment of pro-caspase-9 to oligomerized
	APAF1, thereby selectively suppressing apoptosis initiated via the mitochondrial/cytochrome c
	pathway (PubMed:12773388). {ECO:0000269 PubMed:12773388,
	EC0:0000269 PubMed:22980980, EC0:0000269 PubMed:28935770,
	ECO:0000269 PubMed:29107538, ECO:0000269 PubMed:29158492,
	ECO:0000269 PubMed:30181260}.
Molecular Weight:	9.6 kDa
UniProt:	043504
Pathways:	PI3K-Akt Signaling, Regulation of Cell Size
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	modifications.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN3083158 | 05/07/2024 | Copyright antibodies-online. All rights reserved.

Application Details

During lysate production, the cell wall and other cellular components that are not required for
protein production are removed, leaving only the protein production machinery and the
mitochondria to drive the reaction. During our lysate completion steps, the additional
components needed for protein production (amino acids, cofactors, etc.) are added to produce
something that functions like a cell, but without the constraints of a living system - all that's
needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)