antibodies -online.com





MPZL1 Protein (AA 36-162) (His tag)



Image



Go to Product page

| \sim | |
|--------|-------|
| ()\/⊝ | rview |
| 0 1 0 | VICVV |

| Quantity: | 1 mg |
|-------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target: | MPZL1 |
| Protein Characteristics: | AA 36-162 |
| Origin: | Human |
| Source: | Escherichia coli (E. coli) |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This MPZL1 protein is labelled with His tag. |
| Application: | SDS-PAGE (SDS), ELISA, Western Blotting (WB), Crystallization (Crys) |
| Product Details | |
| Sequence: | SALEVYTPKE IFVANGTQGK LTCKFKSTST TGGLTSVSWS FQPEGADTTV SFFHYSQGQV |
| | YLGNYPPFKD RISWAGDLDK KDASINIENM QFIHNGTYIC DVKNPPDIVV QPGHIRLYVV EKENLPV |
| | Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a |
| | special request, please contact us. |
| Characteristics: | Made in Germany - from design to production - by highly experienced protein experts. Human MPZL1 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade. State-of-the-art algorithm used for plasmid design (Gene synthesis). |
| | This protein is a made to order protein and will be made for the first time for your order. Our |
| | experts in the lab will ensure that you receive a correctly folded protein. |
| | The big advantage of ordering our made-to-order proteins in comparison to ordering custom |
| | made proteins from other companies is that there is no financial obligation in case the protein |
| | |

cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered. The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in bacterial culture:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Endotoxin has not been removed. Please contact us if you require endotoxin removal.

Grade:

Crystallography grade

Target Details

| Target: | MPZL1 |
|-------------------|-------------------------------------------------------------------------------------------------|
| Alternative Name: | MPZL1 (MPZL1 Products) |
| Background: | Cell surface receptor, which is involved in signal transduction processes. Recruits |
| | PTPN11/SHP-2 to the cell membrane and is a putative substrate of PTPN11/SHP-2. Is a major |
| | receptor for concanavalin-A (ConA) and is involved in cellular signaling induced by ConA, which |
| | probably includes Src family tyrosine-protein kinases. Isoform 3 seems to have a dominant |
| | negative role, it blocks tyrosine phosphorylation of MPZL1 induced by ConA. Isoform 1, but not |

Target Details

Expiry Date:

| l arget Details | |
|---------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| | isoform 2 and isoform 3, may be involved in regulation of integrin-mediated cell motility. {ECO:0000269 PubMed:11751924, ECO:0000269 PubMed:12410637}. |
| | {ECO.0000209 Fubivied.11731924, ECO.0000209 Fubivied.12410037}. |
| Molecular Weight: | 15.1 kDa Including tag. |
| UniProt: | 095297 |
| Application Details | |
| Application Notes: | In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though. |
| Comment: | In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. |

Unlimited (if stored properly)



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process