antibodies -online.com





anti-PARP3 antibody (Internal Region)

2 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	PARP3
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This PARP3 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Durnooo:	PARP3
Purpose:	PARPS
Immunogen:	Peptide with sequence YEDYNCTLNQTNIEN, from the internal region of the protein sequence
	according to NP_001003931.2, NP_001003935.2.
Sequence:	YEDYNCTLNQ TNIEN
Isotype:	IgG
130 гуре.	
Specificity:	This antibody is expected to recognize both reported isoforms (NP_001003931.2,
	NP_001003935.2).
Cross-Reactivity:	Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity
	chromatography using the immunizing peptide.


Product Details Grade: Verified **Target Details** Target: PARP3 PARP3 (PARP3 Products) Alternative Name Background: PARP3, poly (ADP-ribose) polymerase family, member 3, ADPRT3, ADPRTL2, ADPRTL3, IRT1, hPARP-3, pADPRT-3, ADP-ribosyltransferase (NAD+, poly (ADP-ribose) polymerase)-like 2, ADPribosyltransferase (NAD+, poly (ADP-ribose) polymerase)-like 3, NAD+ ADP-ribos 10039 Gene ID: NCBI Accession: NP_001003931, NP_001003935 **Application Details Application Notes:** Immunohistochemistry: Paraffin embedded Human Liver and Kidney. Recommended concentration: 3.75 µg/mL. Western Blot: Preliminary experiments in lysates of cell lines Daudi, HeLa and HepG2 gave no specific signal but low background (at antibody concentration up to 1 µg/mL). Peptide ELISA: antibody detection limit dilution 1:2000. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin. Sodium azide Preservative: Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. Minimize freezing and thawing. Handling Advice: -20 °C Storage:

Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated

Storage Comment:


at 4°C for a few weeks and still remain viable.

Images



Immunohistochemistry

Image 1. ABIN308468 (3.75µg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Immunohistochemistry

Image 2. ABIN308468 (3.75µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.