

Datasheet for ABIN3084790

NTAN1 Protein (AA 2-310) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	NTAN1
Protein Characteristics:	AA 2-310
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NTAN1 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

Product Details

Sequence: PLLVEGRRVR LPQSAGDLVR AHPPLEERAR LLRGQSVQQV GPQGLLYVQQ RELAVTSPKD
 GSISILGSDD ATTCHIVLRL HTGNGATCLT HCDGTDTKAE VPLIMNSIKS FSDHAQCGRL
 EVHLVGGFSD DRQLSQKLTH QLLSEFDRQE DDIHLVTLCV TELNDRENE NHFPVIYGIA
 VNIKTAEIYR ASFQDRGPEE QLRAARTLAG GPMISYDAE TEQLRIGPYS WTPFPHVDFW
 LHQDDKQILE NLSTSPLAEP PHFVEHIRST LMFLKKHPSP AHTLFSGNKA LLYKKNEDGL
 WEKISSPGS

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

- | | |
|------------------|--|
| Characteristics: | <ul style="list-style-type: none"> • Made in Germany - from design to production - by highly experienced protein experts. • Human NTAN1 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade. • State-of-the-art algorithm used for plasmid design (Gene synthesis). |
|------------------|--|

Product Details

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in baculovirus infected SF9 insect cells: <ol style="list-style-type: none">1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Protein is endotoxin free.
Grade:	Crystallography grade

Target Details

Target:	NTAN1
Alternative Name:	NTAN1 (NTAN1 Products)
Background:	Side-chain deamidation of N-terminal asparagine residues to aspartate. Required for the

Target Details

ubiquitin-dependent turnover of intracellular proteins that initiate with Met-Asn. These proteins are acetylated on the retained initiator methionine and can subsequently be modified by the removal of N-acetyl methionine by acylaminoacid hydrolase (AAH). Conversion of the resulting N-terminal asparagine to aspartate by PNAD renders the protein susceptible to arginylation, polyubiquitination and degradation as specified by the N-end rule. This enzyme does not act on substrates with internal or C-terminal asparagines and does not act on glutamine residues in any position, nor on acetylated N-terminal peptidyl Asn. {ECO:0000269|PubMed:21375249}.

Molecular Weight: 35.5 kDa Including tag.

UniProt: [Q96AB6](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: Unlimited (if stored properly)