

Datasheet for ABIN3084794

NUDT15 Protein (AA 1-164) (Strep Tag)[Go to Product page](#)**1** Image

Overview

Quantity:	1 mg
Target:	NUDT15
Protein Characteristics:	AA 1-164
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This NUDT15 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Sequence:	<p>MTASAPRGR RPGVGVGVV TSCKHPRCVL LGKRKGSVGA GSFQLPGGHL EFGETWEECA QRETWEEAAL HLKNVHFASV VNSFIEKENY HYVTILMKGE VDVTHDSEPK NVEPEKNESW EWVPWEELPP LDQLFWGLRC LKEQGYDPFK EDLNHLVGYK GNHL</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>
Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none">• Made in Germany - from design to production - by highly experienced protein experts.• Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.• These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).

- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®): 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade

Target Details

Target:	NUDT15
Alternative Name:	NUDT15 (NUDT15 Products)
Background:	<p>Nucleotide triphosphate diphosphatase NUDT15 (EC 3.6.1.9) (MutT homolog 2) (MTH2) (Nucleoside diphosphate-linked moiety X motif 15) (Nudix motif 15) (Nucleoside diphosphate-linked to another moiety X hydrolase 15) (Nudix hydrolase 15),FUNCTION: May catalyze the hydrolysis of nucleoside triphosphates including dGTP, dTTP, dCTP, their oxidized forms like 8-oxo-dGTP and the prodrug thiopurine derivatives 6-thio-dGTP and 6-thio-GTP (PubMed:26238318). Could also catalyze the hydrolysis of some nucleoside diphosphate derivatives (PubMed:22556419, PubMed:26238318). Hydrolyzes oxidized nucleosides triphosphates like 8-oxo-dGTP in vitro, but the specificity and efficiency towards these substrates are low. Therefore, the potential in vivo sanitizing role of this enzyme, that would consist in removing oxidatively damaged forms of nucleosides to prevent their incorporation into DNA, is unclear (PubMed:26238318, PubMed:22556419). Through the hydrolysis of thioguanosine triphosphates may participate in the catabolism of thiopurine drugs (PubMed:26238318, PubMed:25108385). May also have a role in DNA synthesis and cell cycle progression by stabilizing PCNA (PubMed:19419956). Exhibits decapping activity towards dpCoA-capped RNAs in vitro (By similarity). {ECO:0000250 UniProtKB:Q8BG93, ECO:0000269 PubMed:19419956, ECO:0000269 PubMed:22556419, ECO:0000269 PubMed:25108385, ECO:0000269 PubMed:26238318}.</p>
Molecular Weight:	18.6 kDa
UniProt:	Q9NV35

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	<p>ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.</p> <p>During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional</p>

Application Details

components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process