



[Go to Product page](#)

Datasheet for ABIN3085937

## PIP4K2A Protein (AA 2-406) (His tag)

### 1 Image

#### Overview

Quantity:	1 mg
Target:	PIP4K2A
Protein Characteristics:	AA 2-406
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This PIP4K2A protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys)

#### Product Details

Sequence: ATPGNLGSSV LASKTKTKKK HFVAQKVKLF RASDPLLSVL MWGVNHSINE LSHVQIPVML  
MPDDFKAYSK IKVDNHLFNK ENMPSHFKFK EYCPMVFRNL RERFGIDDQD FQNSLTRSAP  
LPNDSQARSG ARFHTSYDKR YIIKTITSED VAEMHNILKK YHQYIVECHG ITLLPQFLGM  
YRLNVDGVEI YVIVTRNVFS HRLSVYRKYD LKGSTVAREA SDKEKAKELP TLKDNDFINE  
GQKIYIDDNN KKVFLKLLK DVEFLAQLKL MDYSLLVGIH DVERAEQEEV ECEENDGEEE  
GESDGTHPVG TPPDSPGNTL NSSPPLAPGE FDPNIDVYGI KCHENSPRKE VYFMAIIDIL  
THYDAKKKAA HAAKTVKHGA GAEISTVNPE QYSKRFLDFI GHILT

**Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.**

- Characteristics:
- Made in Germany - from design to production - by highly experienced protein experts.
  - Human PIP4K2A Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.

## Product Details

---

- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

---

Purification:	Two step purification of proteins expressed in bacterial culture: <ol style="list-style-type: none"><li>1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.</li><li>2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.</li></ol>
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Endotoxin has not been removed. Please contact us if you require endotoxin removal.
Grade:	Crystallography grade

---

## Target Details

---

Target:	PIP4K2A
Alternative Name:	PIP4K2A ( <a href="#">PIP4K2A Products</a> )

---

## Target Details

---

**Background:** Catalyzes the phosphorylation of phosphatidylinositol 5-phosphate (PtdIns5P) on the fourth hydroxyl of the myo-inositol ring, to form phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P<sub>2</sub>). May exert its function by regulating the levels of PtdIns5P, which functions in the cytosol by increasing AKT activity and in the nucleus signals through ING2. May regulate the pool of cytosolic PtdIns5P in response to the activation of tyrosine phosphorylation. May negatively regulate insulin-stimulated glucose uptake by lowering the levels of PtdIns5P. May be involved in thrombopoiesis, and the terminal maturation of megakaryocytes and regulation of their size. {ECO:0000269|PubMed:18364242}.

**Molecular Weight:** 47.0 kDa Including tag.

**UniProt:** [P48426](#)

**Pathways:** [Inositol Metabolic Process](#)

## Application Details

---

**Application Notes:** In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

**Comment:** In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

**Buffer:** 100 mM NaCl, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

**Handling Advice:** Avoid repeated freeze-thaw cycles.

**Storage:** -80 °C

**Storage Comment:** Store at -80°C.

**Expiry Date:** Unlimited (if stored properly)



**Image 1.** „Crystallography Grade“ protein due to multi-step, protein-specific purification process