

## Datasheet for ABIN3086720

# PTGR1 Protein (AA 1-329) (Strep Tag)



## Overview

Quantity:	1 mg
Target:	PTGR1
Protein Characteristics:	AA 1-329
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This PTGR1 protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

#### **Product Details**

#### Sequence:

MVRTKTWTLK KHFVGYPTNS DFELKTAELP PLKNGEVLLE ALFLTVDPYM RVAAKRLKEG
DTMMGQQVAK VVESKNVALP KGTIVLASPG WTTHSISDGK DLEKLLTEWP DTIPLSLALG
TVGMPGLTAY FGLLEICGVK GGETVMVNAA AGAVGSVVGQ IAKLKGCKVV GAVGSDEKVA
YLQKLGFDVV FNYKTVESLE ETLKKASPDG YDCYFDNVGG EFSNTVIGQM KKFGRIAICG
AISTYNRTGP LPPGPPPEIV IYQELRMEAF VVYRWQGDAR QKALKDLLKW VLEGKIQYKE
YIIEGFENMP AAFMGMLKGD NLGKTIVKA

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography

- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

## Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- · The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (AliCE®).
Purity:	> 80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Target Details	
Target:	PTGR1
Alternative Name:	PTGR1 (PTGR1 Products)
Background:	Prostaglandin reductase 1 (PRG-1) (15-oxoprostaglandin 13-reductase) (EC 1.3.1.48)

(Dithiolethione-inducible gene 1 protein) (D3T-inducible gene 1 protein) (DIG-1) (Leukotriene B4 12-hydroxydehydrogenase) (NAD(P)H-dependent alkenal/one oxidoreductase) (EC 1.3.1.74), FUNCTION: NAD(P)H-dependent oxidoreductase involved in metabolic inactivation of pro- and anti-inflammatory eicosanoids: prostaglandins (PG), leukotrienes (LT) and lipoxins (LX) (PubMed:25619643). Catalyzes with high efficiency the reduction of the 13,14 double bond of 15-oxoPGs, including 15-oxo-PGE1, 15-oxo-PGE2, 15-oxo-PGF1-alpha and 15-oxo-PGF2-alpha (PubMed:25619643). Catalyzes with lower efficiency the oxidation of the hydroxyl group at C12 of LTB4 and its derivatives, converting them into biologically less active 12-oxo-LTB4 metabolites (PubMed:25619643) (By similarity). Reduces 15-oxo-LXA4 to 13,14 dihydro-15-oxo-LXA4, enhancing neutrophil recruitment at the inflammatory site (By similarity). May play a role in metabolic detoxification of alkenals and ketones. Reduces alpha, beta-unsaturated alkenals and ketones, particularly those with medium-chain length, showing highest affinity toward (2E)decenal and (3E)-3-nonen-2-one (PubMed:25619643). May inactivate 4-hydroxy-2-nonenal, a cytotoxic lipid constituent of oxidized low-density lipoprotein particles (By similarity). {ECO:0000250|UniProtKB:P97584, ECO:0000250|UniProtKB:Q29073, ECO:0000269|PubMed:25619643}.

Molecular Weight:

35.9 kDa

UniProt:

Q14914

#### **Application Details**

**Application Notes:** 

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment:

ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)