

Datasheet for ABIN3087959

APOBEC3F Protein (AA 1-373) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	APOBEC3F
Protein Characteristics:	AA 1-373
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This APOBEC3F protein is labelled with Strep Tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Brand:	ALICE®
Sequence:	<p>MKPHFRNTVE RMYRDTFSYN FYNRPILSRR NTVWLCYEVK TKGPSRPRLD AKIFRGQVYS QPEHHAEMCF LSWFCGNQLP AYKCFQITWF VSWTPCPDCV AKLAEFLAEH PNVTLTISAA RLYYYWERDY RRALCRLSQA GARVKIMDDE EFAYCWENFV YSEGQPFMPW YKFDDNYAFL HRTLKEILRN PMEAMYPHIF YFHFKNLRKA YGRNESWLCF TMEVVKHHSP VSWKRGVFRN QVDPETHCHA ERCFLSWFCD DILSPNTNVE VTWYTSWSPC PECAGEVAEF LARHSNVNLT IFTARLYYFW DTDYQEGLRS LSQEGASVEI MGYKDFKYCW ENFVYNDDEP FKPWKGLKYN FLFLDSKLQE ILE</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>
Characteristics:	Key Benefits:

Product Details

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).
Purity:	> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).
Grade:	custom-made

Target Details

Target:	APOBEC3F
---------	----------

Target Details

Alternative Name:	APOBEC3F (APOBEC3F Products)
Background:	<p>DNA dC->dU-editing enzyme APOBEC-3F (EC 3.5.4.38) (Apolipoprotein B mRNA-editing enzyme catalytic polypeptide-like 3F) (A3F),FUNCTION: DNA deaminase (cytidine deaminase) which acts as an inhibitor of retrovirus replication and retrotransposon mobility via deaminase-dependent and -independent mechanisms. Exhibits antiviral activity against viruse such as HIV-1 or HIV-2 (PubMed:15141007, PubMed:15152192, PubMed:23001005, PubMed:34774569). After the penetration of retroviral nucleocapsids into target cells of infection and the initiation of reverse transcription, it can induce the conversion of cytosine to uracil in the minus-sense single-strand viral DNA, leading to G-to-A hypermutations in the subsequent plus-strand viral DNA (PubMed:15141007). The resultant detrimental levels of mutations in the proviral genome, along with a deamination-independent mechanism that works prior to the proviral integration, together exert efficient antiretroviral effects in infected target cells. Selectively targets single-stranded DNA and does not deaminate double-stranded DNA or single- or double-stranded RNA. Exhibits antiviral activity also against hepatitis B virus (HBV), equine infectious anemia virus (EIAV), xenotropic MuLV-related virus (XMRV) and simian foamy virus (SFV) and may inhibit the mobility of LTR and non-LTR retrotransposons. May also play a role in the epigenetic regulation of gene expression through the process of active DNA demethylation.</p> <p>{ECO:0000269 PubMed:15141007, ECO:0000269 PubMed:15152192, ECO:0000269 PubMed:16378963, ECO:0000269 PubMed:16527742, ECO:0000269 PubMed:19458006, ECO:0000269 PubMed:20062055, ECO:0000269 PubMed:20219927, ECO:0000269 PubMed:20335265, ECO:0000269 PubMed:21496894, ECO:0000269 PubMed:21835787, ECO:0000269 PubMed:22807680, ECO:0000269 PubMed:22915799, ECO:0000269 PubMed:23001005, ECO:0000269 PubMed:23097438, ECO:0000269 PubMed:23152537, ECO:0000269 PubMed:34774569}.</p>
Molecular Weight:	45.0 kDa
UniProt:	Q8IUX4

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from <i>Nicotiana tabacum</i> c.v.. This contains all the protein expression machinery needed to produce

Application Details

even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months