

Datasheet for ABIN3088013

SLC3A2 Protein (AA 206-630) (His tag)



[Go to Product page](#)

1 Image

Overview

Quantity:	1 mg
Target:	SLC3A2
Protein Characteristics:	AA 206-630
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLC3A2 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence: RAPRCRELPA QKWWHTGALY RIGDLQAFQG HGAGNLAGLK GRLDYLSLKV KGLVLPPIH
KNQKDDVAQT DLLQIDPNFG SKEDFDSLQ SAKKKSIRVI LDLTPNYRGE NSWFSTQVDT
VATKVKDALE FWLQAGVDGF QVRDIENLKD ASSFLAEWQN ITKGFSEDRL LIAGTNSSDL
QQILSLLESN KDLLLTSSYL SDSGSTGEHT KSLVTQYLNA TGNRWCSWSL SQARLLTSFL
PAQLRLYQL MLFTLPGTPV FSYGDEIGLD AAALPGQPME APVMLWDESS FPDIPGAVSA
NMTVKGQSED PGSLLSLFRR LSDQRSKERS LLHGDFHAFS AGPGLFSYIR HWDQNERFLV
VLNFGDVGLS AGLQASDLPA SASLPAKADL LLSTQPGREE GSPLELERLK LEPHEGLLLR FPYAA

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

- Characteristics:
- Made in Germany - from design to production - by highly experienced protein experts.
 - Human SLC3A2 Protein (raised in E. Coli) purified by multi-step, protein-specific process to ensure crystallization grade.

Product Details

- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in bacterial culture: <ol style="list-style-type: none">1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Endotoxin has not been removed. Please contact us if you require endotoxin removal.
Grade:	Crystallography grade

Target Details

Target:	SLC3A2
Alternative Name:	SLC3A2 (SLC3A2 Products)

Target Details

Background: Required for the function of light chain amino-acid transporters. Involved in sodium-independent, high-affinity transport of large neutral amino acids such as phenylalanine, tyrosine, leucine, arginine and tryptophan. Involved in guiding and targeting of LAT1 and LAT2 to the plasma membrane. When associated with SLC7A6 or SLC7A7 acts as an arginine/glutamine exchanger, following an antiport mechanism for amino acid transport, influencing arginine release in exchange for extracellular amino acids. Plays a role in nitric oxide synthesis in human umbilical vein endothelial cells (HUVECs) via transport of L-arginine. Required for normal and neoplastic cell growth. When associated with SLC7A5/LAT1, is also involved in the transport of L-DOPA across the blood-brain barrier, and that of thyroid hormones triiodothyronine (T3) and thyroxine (T4) across the cell membrane in tissues such as placenta. Involved in the uptake of methylmercury (MeHg) when administered as the L-cysteine or D,L-homocysteine complexes, and hence plays a role in metal ion homeostasis and toxicity. When associated with SLC7A5 or SLC7A8, involved in the cellular activity of small molecular weight nitrosothiols, via the stereoselective transport of L-nitrosocysteine (L-CNSO) across the transmembrane. Together with ICAM1, regulates the transport activity LAT2 in polarized intestinal cells, by generating and delivering intracellular signals. When associated with SLC7A5, plays an important role in transporting L-leucine from the circulating blood to the retina across the inner blood-retinal barrier. {ECO:0000269|PubMed:10903140, ECO:0000269|PubMed:11311135, ECO:0000269|PubMed:11389679, ECO:0000269|PubMed:11557028, ECO:0000269|PubMed:11564694, ECO:0000269|PubMed:11742812, ECO:0000269|PubMed:12117417, ECO:0000269|PubMed:12225859, ECO:0000269|PubMed:12716892, ECO:0000269|PubMed:14603368, ECO:0000269|PubMed:15769744, ECO:0000269|PubMed:15980244, ECO:0000269|PubMed:9751058, ECO:0000269|PubMed:9829974, ECO:0000269|PubMed:9878049}.

Molecular Weight: 47.8 kDa Including tag.

UniProt: [P08195](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to

Application Details

increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 100 mM NaCl, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: Unlimited (if stored properly)

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process