# antibodies .- online.com





## APOBEC3G Protein (AA 1-384) (Strep Tag)



**Image** 



Go to Product page

#### Overview

| Quantity:                     | 1 mg  |
|-------------------------------|---|
| Target:                       | APOBEC3G  |
| Protein Characteristics:      | AA 1-384  |
| Origin:                       | Human   |
| Source:                       | Tobacco (Nicotiana tabacum)                       |
| Protein Type:                 | Recombinant                                       |
| Purification tag / Conjugate: | This APOBEC3G protein is labelled with Strep Tag. |
| Application:                  | ELISA, Western Blotting (WB), SDS-PAGE (SDS)      |

#### **Product Details**

Sequence:

MKPHFRNTVE RMYRDTFSYN FYNRPILSRR NTVWLCYEVK TKGPSRPPLD AKIFRGQVYS
ELKYHPEMRF FHWFSKWRKL HRDQEYEVTW YISWSPCTKC TRDMATFLAE DPKVTLTIFV
ARLYYFWDPD YQEALRSLCQ KRDGPRATMK IMNYDEFQHC WSKFVYSQRE LFEPWNNLPK
YYILLHIMLG EILRHSMDPP TFTFNFNNEP WVRGRHETYL CYEVERMHND TWVLLNQRRG
FLCNQAPHKH GFLEGRHAEL CFLDVIPFWK LDLDQDYRVT CFTSWSPCFS CAQEMAKFIS
KNKHVSLCIF TARIYDDQGR CQEGLRTLAE AGAKISIMTY SEFKHCWDTF VDHQGCPFQP
WDGLDEHSQD LSGRLRAILQ NQEN

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- · Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
  protein production are removed, leaving only the protein production machinery and the
  mitochondria to drive the reaction. During our lysate completion steps, the additional
  components needed for protein production (amino acids, cofactors, etc.) are added to
  produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- · We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

#### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

- 1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

### **Product Details**

| Purity:          | >80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot. |
|------------------|--|
| Endotoxin Level: | Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)                                    |
| Grade:           | Crystallography grade  |

| Grade:            | Crystallography grade  |
|-------------------|--|
| Target Details    |  |
| Target:           | APOBEC3G   |
| Alternative Name: | APOBEC3G (APOBEC3G Products)   |
| Background:       | DNA dC->dU-editing enzyme APOBEC-3G (EC 3.5.4.38) (APOBEC-related cytidine deaminase)                  |
|                   | (APOBEC-related protein) (ARCD) (APOBEC-related protein 9) (ARP-9) (CEM-15) (CEM15)                    |
|                   | (Deoxycytidine deaminase) (A3G),FUNCTION: DNA deaminase (cytidine deaminase) which acts                |
|                   | as an inhibitor of retrovirus replication and retrotransposon mobility via deaminase-dependent         |
|                   | and -independent mechanisms. Exhibits potent antiviral activity against Vif-deficient HIV-1.           |
|                   | After the penetration of retroviral nucleocapsids into target cells of infection and the initiation of |
|                   | reverse transcription, it can induce the conversion of cytosine to uracil in the minus-sense           |
|                   | single-strand viral DNA, leading to G-to-A hypermutations in the subsequent plus-strand viral          |
|                   | DNA. The resultant detrimental levels of mutations in the proviral genome, along with a                |
|                   | deamination-independent mechanism that works prior to the proviral integration, together exert         |
|                   | efficient antiretroviral effects in infected target cells. Selectively targets single-stranded DNA     |
|                   | and does not deaminate double-stranded DNA or single- or double-stranded RNA. Exhibits                 |
|                   | antiviral activity also against simian immunodeficiency viruses (SIVs), hepatitis B virus (HBV),       |
|                   | equine infectious anemia virus (EIAV), xenotropic MuLV-related virus (XMRV) and simian foamy           |
|                   | virus (SFV). May inhibit the mobility of LTR and non-LTR retrotransposons.                             |
|                   | {ECO:0000269 PubMed:12167863, ECO:0000269 PubMed:12808465,   |
|                   | ECO:0000269 PubMed:12808466, ECO:0000269 PubMed:12809610,  |
|                   | ECO:0000269 PubMed:12859895, ECO:0000269 PubMed:12970355,  |
|                   | ECO:0000269 PubMed:14528300, ECO:0000269 PubMed:14557625,  |
|                   | ECO:0000269 PubMed:15031497, ECO:0000269 PubMed:16378963,  |
|                   | ECO:0000269 PubMed:16527742, ECO:0000269 PubMed:18288108,  |
|                   | ECO:0000269 PubMed:19458006, ECO:0000269 PubMed:20219927,  |
|                   | ECO:0000269 PubMed:20335265, ECO:0000269 PubMed:21123384,  |
|                   | ECO:0000269 PubMed:21835787, ECO:0000269 PubMed:22791714,  |
|                   | ECO:0000269 PubMed:22807680, ECO:0000269 PubMed:22915799,  |
|                   | ECO:0000269 PubMed:23097438, ECO:0000269 PubMed:23152537}.   |

## Target Details

| Target Details      |   |
|---------------------|---|
| Molecular Weight:   | 46.4 kDa  |
| UniProt:            | Q9HC16  |
| Application Details |   |
| Application Notes:  | In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.  |
| Comment:            | ALICE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.   |
|                     | During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce |
|                     | something that functions like a cell, but without the constraints of a living system - all that's   |

Restrictions:

For Research Use only

## Handling

| Format:          | Liquid   |
|------------------|--|
| Buffer:          | The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us. |
| Handling Advice: | Avoid repeated freeze-thaw cycles.   |
| Storage:         | -80 °C   |
| Storage Comment: | Store at -80°C.  |
| Expiry Date:     | Unlimited (if stored properly)   |

needed is the DNA that codes for the desired protein!



**Image 1.** "Crystallography Grade" protein due to multi-step, protein-specific purification process