
antibodies .- online.com





ADAM15 Protein (AA 207-696) (His tag)



Overview

| Quantity: | 1 mg |
|-------------------------------|--|
| Target: | ADAM15 |
| Protein Characteristics: | AA 207-696 |
| Origin: | Human |
| Source: | Insect Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This ADAM15 protein is labelled with His tag. |
| Application: | ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys) |

Product Details

Sequence:

DVVTETKTVE LVIVADHSEA QKYRDFQHLL NRTLEVALLL DTFFRPLNVR VALVGLEAWT
QRDLVEISPN PAVTLENFLH WRRAHLLPRL PHDSAQLVTG TSFSGPTVGM AIQNSICSPD
FSGGVNMDHS TSILGVASSI AHELGHSLGL DHDLPGNSCP CPGPAPAKTC IMEASTDFLP
GLNFSNCSRR ALEKALLDGM GSCLFERLPS LPPMAAFCGN MFVEPGEQCD CGFLDDCVDP
CCDSLTCQLR PGAQCASDGP CCQNCQLRPS GWQCRPTRGD CDLPEFCPGD SSQCPPDVSL
GDGEPCAGGQ AVCMHGRCAS YAQQCQSLWG PGAQPAAPLC LQTANTRGNA FGSCGRNPSG
SYVSCTPRDA ICGQLQCQTG RTQPLLGSIR DLLWETIDVN GTELNCSWVH LDLGSDVAQP
LLTLPGTACG PGLVCIDHRC QRVDLLGAQE CRSKCHGHGV CDSNRHCYCE EGWAPPDCTT
QLKATSSLTT

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human ADAM15 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free.

Grade:

Crystallography grade

Target Details

| Target: | ADAM15 |
|---------------------|--|
| Alternative Name: | ADAM15 (ADAM15 Products) |
| Background: | Active metalloproteinase with gelatinolytic and collagenolytic activity. Plays a role in the wound |
| | healing process. Mediates both heterotypic intraepithelial cell/T-cell interactions and homotypic |
| | T-cell aggregation. Inhibits beta-1 integrin-mediated cell adhesion and migration of airway |
| | smooth muscle cells. Suppresses cell motility on or towards fibronectin possibly by driving |
| | alpha-v/beta-1 integrin (ITAGV-ITGB1) cell surface expression via ERK1/2 inactivation. Cleaves |
| | E-cadherin in response to growth factor deprivation. Plays a role in glomerular cell migration. |
| | Plays a role in pathological neovascularization. May play a role in cartilage remodeling. May be |
| | proteolytically processed, during sperm epididymal maturation and the acrosome reaction. May |
| | play a role in sperm-egg binding through its disintegrin domain. |
| | {ECO:0000269 PubMed:12091380, ECO:0000269 PubMed:15358598, |
| | ECO:0000269 PubMed:15818704, ECO:0000269 PubMed:17416588, |
| | ECO:0000269 PubMed:17575078, ECO:0000269 PubMed:18387333, |
| | ECO:0000269 PubMed:18434311}. |
| Molecular Weight: | 53.3 kDa Including tag. |
| UniProt: | Q13444 |
| Application Details | |
| Application Notes: | In addition to the applications listed above we expect the protein to work for functional studies |
| | as well. As the protein has not been tested for functional studies yet we cannot offer a |
| | guarantee though. |
| Comment: | In cases in which it is highly likely that the recombinant protein with the default tag will be |
| | insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to |
| | increase solubility. We will discuss all possible options with you in detail to assure that you |
| | receive your protein of interest. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| | |
| Buffer: | 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer. |

Handling

| Storage: | -80 °C |
|------------------|--------------------------------|
| Storage Comment: | Store at -80°C. |
| Expiry Date: | Unlimited (if stored properly) |