antibodies .- online.com





ACOX1 Protein (AA 1-660) (Strep Tag)



Image



Go to Product page

Overview

Quantity:	1 mg
Target:	ACOX1
Protein Characteristics:	AA 1-660
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This ACOX1 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

Product Details

Sequence:

MNPDLRRERD SASFNPELLT HILDGSPEKT RRRREIENMI LNDPDFQHED LNFLTRSQRY
EVAVRKSAIM VKKMREFGIA DPDEIMWFKK LHLVNFVEPV GLNYSMFIPT LLNQGTTAQK
EKWLLSSKGL QIIGTYAQTE MGHGTHLRGL ETTATYDPET QEFILNSPTV TSIKWWPGGL
GKTSNHAIVL AQLITKGKCY GLHAFIVPIR EIGTHKPLPG ITVGDIGPKF GYDEIDNGYL
KMDNHRIPRE NMLMKYAQVK PDGTYVKPLS NKLTYGTMVF VRSFLVGEAA RALSKACTIA
IRYSAVRHQS EIKPGEPEPQ ILDFQTQQYK LFPLLATAYA FQFVGAYMKE TYHRINEGIG
QGDLSELPEL HALTAGLKAF TSWTANTGIE ACRMACGGHG YSHCSGLPNI YVNFTPSCTF
EGENTVMMLQ TARFLMKSYD QVHSGKLVCG MVSYLNDLPS QRIQPQQVAV WPTMVDINSP
ESLTEAYKLR AARLVEIAAK NLQKEVIHRK SKEVAWNLTS VDLVRASEAH CHYVVVKLFS
EKLLKIQDKA IQAVLRSLCL LYSLYGISQN AGDFLQGSIM TEPQITQVNQ RVKELLTLIR
SDAVALVDAF DFQDVTLGSV LGRYDGNVYE NLFEWAKNSP LNKAEVHESY KHLKSLQSKL

Sequence without tag. The proposed Strep-Tag is based on experience s with the expression

system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany from design to production by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- · State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag

UniProt:

Pathways:

Q15067

Product Details			
	capture material. Eluate fractions are analyzed by SDS-PAGE. 2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.		
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.		
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)		
Grade:	Crystallography grade		
Target Details			
Target:	ACOX1		
Alternative Name:	ACOX1 (ACOX1 Products)		
Background:	Peroxisomal acyl-coenzyme A oxidase 1 (AOX) (EC 1.3.3.6) (Palmitoyl-CoA oxidase)		
	(Peroxisomal fatty acyl-CoA oxidase) (Straight-chain acyl-CoA oxidase) (SCOX) [Cleaved into:		
	Peroxisomal acyl-CoA oxidase 1, A chain, Peroxisomal acyl-CoA oxidase 1, B chain,		
	Peroxisomal acyl-CoA oxidase 1, C chain], FUNCTION: Involved in the initial and rate-limiting		
	step of peroxisomal beta-oxidation of straight-chain saturated and unsaturated very-long-chair		
	fatty acids (PubMed:7876265, PubMed:15060085, PubMed:17458872, PubMed:17603022,		
	PubMed:32169171, PubMed:33234382). Catalyzes the desaturation of fatty acyl-CoAs such as		
	palmitoyl-CoA (hexadecanoyl-CoA) to 2-trans-enoyl-CoAs ((2E)-enoyl-CoAs) such as (2E)-		
	hexadecenoyl-CoA, and donates electrons directly to molecular oxygen (O(2)), thereby		
	producing hydrogen peroxide (H(2)O(2)) (PubMed:7876265, PubMed:17458872,		
	PubMed:17603022). {ECO:0000269 PubMed:15060085, ECO:0000269 PubMed:17458872,		
	ECO:0000269 PubMed:17603022, ECO:0000269 PubMed:32169171,		
	ECO:0000269 PubMed:33234382, ECO:0000269 PubMed:7876265}., FUNCTION: [Isoform 1]:		
	Shows highest activity against medium-chain fatty acyl-CoAs. Shows optimum activity with a		
	chain length of 10 carbons (decanoyl-CoA) in vitro. {ECO:0000269 PubMed:17603022}.,		
	FUNCTION: [Isoform 2]: Is active against a much broader range of substrates and shows		
	activity towards long-chain fatty acyl-CoAs. {ECO:0000269 PubMed:17603022}.		
Molecular Weight:	74.4 kDa		

Regulation of Lipid Metabolism by PPARalpha, Monocarboxylic Acid Catabolic Process

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.	
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the	
	mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	Unlimited (if stored properly)	



Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process