antibodies .- online.com





ANAPC2 Protein (AA 1-822) (His tag)





Overview

Quantity:	1 mg
Target:	ANAPC2
Protein Characteristics:	AA 1-822
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ANAPC2 protein is labelled with His tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

Product Details

Sequence:

MAAAVVVAEG DSDSRPGQEL LVAWNTVSTG LVPPAALGLV SSRTSGAVPP KEEELRAAVE VLRGHGLHSV LEEWFVEVLQ NDLQANISPE FWNAISQCEN SADEPQCLLL LLDAFGLLES RLDPYLRSLE LLEKWTRLGL LMGTGAQGLR EEVHTMLRGV LFFSTPRTFQ EMIQRLYGCF LRVYMQSKRK GEGGTDPELE GELDSRYARR RYYRLLQSPL CAGCSSDKQQ CWCRQALEQF HQLSQVLHRL SLLERVSAEA VTTTLHQVTR ERMEDRCRGE YERSFLREFH KWIERVVGWL GKVFLQDGPA RPASPEAGNT LRRWRCHVQR FFYRIYASLR IEELFSIVRD FPDSRPAIED LKYCLERTDQ RQQLLVSLKA ALETRLLHPG VNTCDIITLY ISAIKALRVL DPSMVILEVA CEPIRRYLRT REDTVRQIVA GLTGDSDGTG DLAVELSKTD PASLETGQDS EDDSGEPEDW VPDPVDADPG KSSSKRRSSD IISLLVSIYG SKDLFINEYR SLLADRLLHQ FSFSPEREIR NVELLKLRFG EAPMHFCEVM LKDMADSRRI NANIREEDEK RPAEEQPPFG VYAVILSSEF WPPFKDEKLE VPEDIRAALE AYCKKYEQLK AMRTLSWKHT LGLVTMDVEL ADRTLSVAVT PVQAVILLYF QDQASWTLEE LSKAVKMPVA LLRRRMSVWL QQGVLREEPP GTFSVIEEER

PQDRDNMVLI DSDDESDSGM ASQADQKEEE LLLFWTYIQA MLTNLESLSL DRIYNMLRMF VVTGPALAEI DLQELQGYLQ KKVRDQQLVY SAGVYRLPKN CS

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany from design to production by highly experienced protein experts.
- Human ANAPC2 Protein (raised in Insect Cells) purified by multi-step, protein-specific process to ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receival of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm.

The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:

Two step purification of proteins expressed in baculovirus infected SF9 insect cells:

- 1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.
- Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity:

>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility:

0.22 µm filtered

Endotoxin Level:

Protein is endotoxin free

Product Details	
Grade:	Crystallography grade
Target Details	
Target:	ANAPC2
Alternative Name:	ANAPC2 (ANAPC2 Products)
Background: Molecular Weight:	Together with the RING-H2 protein ANAPC11, constitutes the catalytic component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains. The CDC20-APC/C complex positively regulates the formation of synaptic vesicle clustering at active zone to the presynaptic membrane in postmitotic neurons. CDC20-APC/C-induced degradation of NEUROD2 drives presynaptic differentiation. {ECO:0000269 PubMed:11739784, ECO:0000269 PubMed:18485873}.
UniProt:	Q9UJX6
Pathways:	Regulation of Cell Size
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling

Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process