antibodies .- online.com





AP4S1 Protein (AA 1-144) (Strep Tag)



Image



Overview

Quantity:	1 mg
Target:	AP4S1 (AP4s1)
Protein Characteristics:	AA 1-144
Origin:	Human
Source:	Tobacco (Nicotiana tabacum)
Protein Type:	Recombinant
Purification tag / Conjugate:	This AP4S1 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Purification tag / Conjugate:	This AP4S1 protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)
Product Details	
Sequence:	MIKFFLMVNK QGQTRLSKYY EHVDINKRTL LETEVIKSCL SRSNEQCSFI EYKDFKLIYR
	QYAALFIVVG VNDTENEMAI YEFIHNFVEV LDEYFSRVSE LDIMFNLDKV HIILDEMVLN
	GCIVETNRAR ILAPLLILDK MSES
	Sequence without tag. The proposed Strep-Tag is based on experience s with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:
	Made in Germany - from design to production - by highly experienced protein experts.
	• Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.

reported (not tested by us and not guaranteed).

· These proteins are normally active (enzymatically functional) as our customers have

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require posttranslational modifications.
- During lysate production, the cell wall and other cellular components that are not required for
 protein production are removed, leaving only the protein production machinery and the
 mitochondria to drive the reaction. During our lysate completion steps, the additional
 components needed for protein production (amino acids, cofactors, etc.) are added to
 produce something that functions like a cell, but without the constraints of a living system all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in Almost Living Cell-Free Expression System
	(ALiCE®):
	1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
	Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
Purity:	>80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Endotoxin Level:	Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)
Grade:	Crystallography grade

Target Details

Target:	AP4S1 (AP4s1)
Alternative Name:	AP4S1 (AP4s1 Products)
Background:	AP-4 complex subunit sigma-1 (AP-4 adaptor complex subunit sigma-1) (Adaptor-related
	protein complex 4 subunit sigma-1) (Sigma-1 subunit of AP-4) (Sigma-4-adaptin) (Sigma4-
	adaptin), FUNCTION: Component of the adaptor protein complex 4 (AP-4). Adaptor protein
	complexes are vesicle coat components involved both in vesicle formation and cargo selection
	They control the vesicular transport of proteins in different trafficking pathways
	(PubMed:10066790, PubMed:10436028). AP-4 forms a non clathrin-associated coat on vesicles
	departing the trans-Golgi network (TGN) and may be involved in the targeting of proteins from
	the trans-Golgi network (TGN) to the endosomal-lysosomal system. It is also involved in protein
	sorting to the basolateral membrane in epithelial cells and the proper asymmetric localization
	of somatodendritic proteins in neurons. AP-4 is involved in the recognition and binding of
	tyrosine-based sorting signals found in the cytoplasmic part of cargos, but may also recognize
	other types of sorting signal (Probable). {ECO:0000269 PubMed:10066790,
	ECO:0000269 PubMed:10436028, ECO:0000305 PubMed:10066790,
	ECO:0000305 PubMed:10436028}.
Molecular Weight:	17.0 kDa
UniProt:	Q9Y587
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from
	Nicotiana tabacum c.v This contains all the protein expression machinery needed to produce
	even the most difficult-to-express proteins, including those that require post-translational
	even the most difficult-to-express proteins, including those that require post-translational modifications.
	modifications.
	modifications. During lysate production, the cell wall and other cellular components that are not required for
	modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the
	modifications. During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

Images




Image 1. "Crystallography Grade" protein due to multi-step, protein-specific purification process