

Datasheet for ABIN3088900

APLF Protein (AA 1-511) (Strep Tag)



[Go to Product page](#)

Overview

Quantity:	250 µg
Target:	APLF
Protein Characteristics:	AA 1-511
Origin:	Human
Source:	Cell-free protein synthesis (CFPS)
Protein Type:	Recombinant
Purification tag / Conjugate:	This APLF protein is labelled with Strep Tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Brand:	AliCE®
Sequence:	<p>MSGGFELQPR DGGPRVALAP GETVIGRGPL LGITDKRVSR RHAILEVAGG QLRIKPIHTN PCFYQSSEKS QLLPLKPNLW CYLNP GDSFS LLVDKYIFRI LSIPSEVEMQ CTLRNSQVLD EDNILNETPK SPVINLPHE T GASQLEGST EIAKTQMTPT NSVSFLGENR DCNKQQPILA ERKRILPTWM LAEHLSDQNL SVPAISGGNV IQGSGKEEIC KDKSQLNTTQ QGRRQLISSG SSENTSAEQD TGEECKNTDQ EESTISSKEM PQSFSAITLS NTEMNNIKTN AQRNKLPIEE LGKVSKHKIA TKRTPHKEDE AMSCSENCSS AQGDSLQDES QGSHSESSSN PSNPETLHAK ATDSVLQGSE GNKVKRTSCM YGANCYRKNP VHFQHF SHPG DSDYGGVQIV GQDETDDRPE CPYGPSCYRK NPQHKIEYRH NTL PVRNVLD EDNDNVGQPN EYDLNDSFLD DEEEDYEPTD EDSDWEPGKE DEEKEDVEEL LKEAKRFMKR K</p> <p>Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you</p>

have a special request, please contact us.

Characteristics:

Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified in one-step affinity chromatography
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured against its specific reference buffer.
- We use the ExPASy's ProtParam tool to determine the absorption coefficient of each protein.

Purification:

One-step Strep-tag purification of proteins expressed in Almost Living Cell-Free Expression System (ALiCE®).

Purity:

> 70-80 % as determined by SDS PAGE, Western Blot and analytical SEC (HPLC).

Grade:

custom-made

Target Details

Target:	APLF
Alternative Name:	APLF (APLF Products)
Background:	<p>Aprataxin and PNK-like factor (EC 3.1.-.-) (Apurinic-apyrimidinic endonuclease APLF) (PNK and APTX-like FHA domain-containing protein) (XRCC1-interacting protein 1),FUNCTION: Histone chaperone involved in single-strand and double-strand DNA break repair (PubMed:17353262, PubMed:17396150, PubMed:21211721, PubMed:21211722, PubMed:30104678, PubMed:29905837). Recruited to sites of DNA damage through interaction with branched poly-ADP-ribose chains, a polymeric post-translational modification synthesized transiently at sites of chromosomal damage to accelerate DNA strand break repair reactions (PubMed:17353262, PubMed:17396150, PubMed:21211721, PubMed:30104678). Following recruitment to DNA damage sites, acts as a histone chaperone that mediates histone eviction during DNA repair and promotes recruitment of histone variant MACROH2A1 (PubMed:21211722, PubMed:30104678, PubMed:29905837). Also has a nuclease activity: displays apurinic-apyrimidinic (AP) endonuclease and 3'-5' exonuclease activities in vitro (PubMed:17353262, PubMed:17396150). Also able to introduce nicks at hydroxyuracil and other types of pyrimidine base damage (PubMed:17353262, PubMed:17396150). Together with PARP3, promotes the retention of the LIG4-XRCC4 complex on chromatin and accelerate DNA ligation during non-homologous end-joining (NHEJ) (PubMed:21211721, PubMed:23689425). Also acts as a negative regulator of cell pluripotency by promoting histone exchange (By similarity). Required for the embryo implantation during the epithelial to mesenchymal transition in females (By similarity). {ECO:0000250 UniProtKB:Q9D842, ECO:0000269 PubMed:17353262, ECO:0000269 PubMed:17396150, ECO:0000269 PubMed:21211721, ECO:0000269 PubMed:21211722, ECO:0000269 PubMed:23689425, ECO:0000269 PubMed:29905837, ECO:0000269 PubMed:30104678}.</p>
Molecular Weight:	57.0 kDa
UniProt:	Q8IW19
Pathways:	DNA Damage Repair

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from

Application Details

Nicotiana tabacum c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. Standard Storage Buffer: PBS pH 7.4, 10 % Glycerol Might differ depending on protein.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months